

Restricted food intake a predictor of increased suicide attempts in Body Dysmorphic Disorder patients

June 5 2012

Rhode Island Hospital and Auburn University researchers found a link between restrictive food intake, or excessive dieting, and an increase in suicide attempts in people with Body Dysmorphic Disorder (BDD). The study focused on the acquired capability of suicide, which is one component of Joiner's (2005) interpersonal-psychological theory of suicide and is comprised of physical pain tolerance and lowered fear of death. The paper is published in the journal *Suicide and Life-Threatening Behavior*, and is now available online in advance of print.

BDD is a common, often severe, and under-recognized [body image](#) disorder. People with BDD experience distressing or impairing preoccupations with perceived flaws in their appearance and are obsessed with the belief that something is wrong with how they look, when in reality they look normal. More than 75 percent of people with BDD feel life is not worth living or think about suicide in their lifetime, and approximately 25 percent have a history of a suicide attempt.

Authors Tracy K. Witte, Ph.D., of Auburn University, Elizabeth R. Didie, Ph.D., a psychologist in the department of psychiatry at Rhode Island Hospital, and Katharine A. Phillips, M.D., director of the [Body Dysmorphic Disorder](#) Program at Rhode Island Hospital examined the association of [suicide attempts](#) with physically painful BDD-related behaviors, including restrictive food intake, excessive exercise, BDD-related [cosmetic surgery](#), compulsive skin picking, and physical self-

mutilation.

The study found that BDD-related restrictive food intake was associated with more than double the number of suicide attempts, but was not associated with suicide ideation; and that those with a history of BDD-related [excessive exercise](#) had less than half the number of suicide attempts as those without such a history. The study also found that none of the other variables indicating exposure to painful and provocative experiences, such as BDD-related cosmetic surgery and compulsive skin picking, were significant predictors of suicide attempts.

Because restriction of food intake can be physically painful, the researchers theorize that a person who is capable of enduring the [physical discomfort](#) of caloric restriction may be more capable of enduring the physical discomfort required in order to inflict self-harm. They theorize that severe restriction of food intake that results in long-standing physical discomfort would predict capability for suicide, whereas more moderate dieting behaviors would have less of a relationship (if any).

"Significantly limiting [food intake](#) can be physically painful," Phillips said. "It goes against our natural instincts to feed our bodies and respond to the physical pain that comes with extreme hunger. The results of this study suggest the importance of assessing individuals with BDD for restrictive eating behaviors to identify suicide risk, even if they have not previously been diagnosed with an eating disorder."

The study included interviews with 200 individuals (68.5 percent women) between the ages of 14 and 64 who had a lifetime diagnosis of BDD. The main criterion variable was the participants' number of past suicide attempts, which ranged from 0 to 25 in the study group. Additionally, 78 percent of the study group had a history of suicide-related ideation. The study only included examination of suicide

attempts, not deaths by suicide.

"While some of the other BDD-related behaviors may seem outwardly more painful – such as undergoing repeated cosmetic procedures, or compulsive skin picking, the level of pain associated with excessive dieting could significantly increase a person's [pain tolerance](#)," Didie said. "This study suggests that those who are capable of enduring such physical discomfort and pain from restrictive eating also may be capable of enduring the physical discomfort required to inflict self harm."

Provided by Lifespan

Citation: Restricted food intake a predictor of increased suicide attempts in Body Dysmorphic Disorder patients (2012, June 5) retrieved 30 April 2024 from <https://medicalxpress.com/news/2012-06-restricted-food-intake-predictor-suicide.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--