


Ringer's acetate better for patients with severe sepsis

June 28 2012



Fluid resuscitation with hydroxyethyl starch 130/0.4 for patients with severe sepsis leads to an increased risk of death at day 90 and an increased likelihood of requiring renal-replacement therapy, compared with Ringer's acetate, according to a study published online June 27 in the *New England Journal of Medicine*.

(HealthDay) -- Fluid resuscitation with hydroxyethyl starch (HES) 130/0.4 for patients with severe sepsis leads to an increased risk of death at day 90 and an increased likelihood of requiring renal-replacement therapy, compared with Ringer's acetate, according to a study published online June 27 in the *New England Journal of Medicine*.

Anders Perner, M.D., Ph.D., from Rigshospitalet in Copenhagen, Denmark, and colleagues randomly assigned 798 participants with severe sepsis participating in a multicenter trial to fluid resuscitation in the [intensive care unit](#) with 6 percent HES 130/0.4 or Ringer's acetate (up to 33 mL per kilogram of ideal body weight per day).

The researchers found that, at 90 days following randomization, 201 of 398 patients assigned to HES 130/0.4 had died, compared to 172 of 400 patients assigned to Ringer's acetate (relative risk [RR], 1.17; 95 percent confidence interval [CI], 1.01 to 1.36), and one patient in each group had end-stage [kidney failure](#). In the HES 130/0.4 group, 87 patients (22 percent) were treated with renal replacement therapy in the 90-day period, compared to 65 patients (16 percent) in the Ringer's acetate group (RR, 1.35; 95 percent CI, 1.01 to 1.80), and 38 and 25 patients, respectively, had severe bleeding (RR, 1.52; 95 percent CI, 0.94 to 2.48).

"Patients with severe [sepsis](#) assigned to fluid resuscitation with HES 130/0.4 had an increased risk of death at day 90 and were more likely to require renal-replacement therapy, as compared with those receiving Ringer's acetate," the authors write.

Perner disclosed receiving grant support from Fresenius Kabi.

More information: [Abstract](#)
[Full Text](#)

Copyright © 2012 [HealthDay](#). All rights reserved.

Citation: Ringer's acetate better for patients with severe sepsis (2012, June 28) retrieved 26 April 2024 from <https://medicalxpress.com/news/2012-06-ringer-acetate-patients-severe-sepsis.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--