

Receiving chemotherapy following removal of type of cancer near pancreas may improve survival

July 10 2012

Patients who had surgery for periampullary cancer (a variety of types of cancer that are located in and near the head of the pancreas, including an area called the ampulla where the bile duct joins up with the pancreatic duct to empty their secretions into the upper small intestine) and received chemotherapy had a statistically significant survival benefit, compared to patients who did not receive chemotherapy, after adjusting for prognostic variables, according to a study in the July 11 issue of *JAMA*.

Periampullary carcinomas arise from the head of the pancreas. "The clinical presentation is similar to that of pancreatic [ductal adenocarcinoma](#), and together they represent a major cause of death. Around 80 percent of periampullary adenocarcinomas are resectable and thus comprise around 30 percent to 40 percent of all resections for cancers in the head of the pancreas," according to background information in the article. Although chemotherapy after surgery has been shown to have a [survival](#) benefit for pancreatic cancer, there have been no [randomized trials](#) for periampullary adenocarcinomas.

John P. Neoptolemos, M.D., of the University of Liverpool, England, and colleagues conducted a study to determine whether adjuvant chemotherapy (fluorouracil or gemcitabine) provides improved overall survival following resection. The [randomized controlled trial](#) (July 2000-May 2008) was conducted in 100 centers in Europe, Australia,

Japan, and Canada. Of the 428 patients included in the primary analysis, 297 had ampullary, 96 had bile duct, and 35 had other cancers. One hundred forty-four patients were assigned to the observation group, 143 patients to receive folinic acid via intravenous bolus injection followed by fluorouracil via intravenous bolus injection administered 1 to 5 days every 28 days, and 141 patients to receive intravenous infusion of gemcitabine once a week for 3 of every 4 weeks for 6 months.

Two hundred forty-four patients (57 percent) had died at the time of analysis, 88 (61 percent) in the observation group, 83 (58 percent) in the fluorouracil plus folinic acid group, and 73 (52 percent) in the gemcitabine group. For the primary analysis, in the observation group, the median (midpoint) survival was 35.2 months and in the chemotherapy group 43.1 months.

The unadjusted primary analysis of the primary outcome of survival did not demonstrate a significant benefit for adjuvant chemotherapy.

"Multivariate analysis, correcting for prognostic variables, found a statistically significant [survival benefit](#) to chemotherapy and specifically for [gemcitabine](#) compared with observation, notwithstanding the better safety profile compared with fluorouracil plus folinic acid, but these results should be considered hypothesis generating. There were different survival outcomes by tumor type, although age, poorly differentiated tumor grade, and lymph node involvement were also independent survival factors," the authors write.

"Although this study found support for the use adjuvant chemotherapy to improve survival in patients with periampullary cancers, this effect was modest, indicating a need for further improvements and warranting the testing of combination chemotherapies."

More information: *JAMA*. 2012;308[2]:147-156.

Provided by JAMA and Archives Journals

Citation: Receiving chemotherapy following removal of type of cancer near pancreas may improve survival (2012, July 10) retrieved 23 April 2024 from <https://medicalxpress.com/news/2012-07-chemotherapy-cancer-pancreas-survival.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.