

Decreasing cancer risk associated with inflammatory bowel disease

July 9 2012

Inflammatory bowel disease is caused by chronic inflammation , which leads to damage of the intestinal epithelium.

Patients with [inflammatory bowel disease](#) have an elevated risk for developing colorectal cancer because of this [chronic inflammation](#).

In an effort to develop strategies to break the cycle of inflammation, Dr. Brent Polk and colleagues at the University of Southern California in Los Angeles examined two mouse models of colorectal cancer.

Their work shows that inactivating a key receptor, known as epidermal growth factor receptor, increases the frequency and severity of colorectal tumors.

Though epidermal growth factor has well-defined roles in promoting tumor growth, the Polk team now finds that epidermal growth factor receptor can ameliorate the response to chronic inflammation and reduces tumor development by fine-tuning inflammation and the generation of intestinal epithelium.

Their results suggest that promoting epidermal growth factor activity in patients with inflammatory bowel disease could decrease long-term [cancer risk](#) by reducing inflammation.

More information: View this article at:
[www.jci.org/articles/view/6288 ... ab673ae8215aaf449677](http://www.jci.org/articles/view/6288...ab673ae8215aaf449677)

Provided by Journal of Clinical Investigation

Citation: Decreasing cancer risk associated with inflammatory bowel disease (2012, July 9)
retrieved 4 May 2024 from

<https://medicalxpress.com/news/2012-07-decreasing-cancer-inflammatory-bowel-disease.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.