

# Heart CT scans may help emergency room personnel more quickly assess patients with chest pain

July 25 2012

---

Adding computed tomography (CT) scans to standard screening procedures may help emergency room staff more rapidly determine which patients complaining of chest pain are having a heart attack or may soon have a heart attack, and which patients can be safely discharged, according to a study funded by the National Heart, Lung, and Blood Institute (NHLBI), part of the National Institutes of Health.

Researchers in the study focused on a condition known as acute coronary syndrome, which includes heart attacks and [unstable angina \(chest pain\)](#), a condition that often progresses to a [heart attack](#). This syndrome occurs when narrowed or blocked coronary arteries prevent oxygen-rich blood from reaching the [heart muscle](#). Since chest pain has many causes, patients are often unnecessarily admitted to the hospital before it is determined that their chest pain is not due to acute coronary syndrome or other serious conditions.

[CT angiography](#) is a type of heart X-ray exam using a device that creates pictures of the coronary arteries, allowing physicians to see whether arteries have major [blockages](#).

"Quickly figuring out which emergency room patients have acute coronary syndrome and which patients can go home because they don't have a serious condition is a significant challenge for U.S. hospitals," said Susan B. Shurin, M.D., acting director of the NHLBI. "Identifying

the underlying cause of chest pain more quickly with CT scans could allow [medical care providers](#) to better allocate limited resources to the patients who are most in need of treatment."

The study results suggest that CT scans allow hospitals to send many patients with chest pain home sooner without compromising their safety. The average length of hospital stay was 23.2 hours for those who underwent CT scans, compared to 30.7 hours for those who underwent standard screening procedures alone. Half of the patients who received a CT scan were discharged in 8.6 hours or less. In contrast, half of the patients in the standard evaluation group were sent home in 26.7 hours or less.

Even with shorter hospital stays in the group that received CT scans, the researchers did not miss any cases of acute coronary syndrome among those participants. After 28 days of follow-up, there was no significant difference in serious cardiovascular events between the two groups.

The study, which appears in the July 26 *New England Journal of Medicine*, was part of an NHLBI-funded program called Rule Out Myocardial Infarction/Ischemia Using Computer-Assisted Tomography (ROMICAT-II).

Led by investigators at Massachusetts General Hospital in Boston, the researchers randomized participants with suspected acute coronary syndrome to receive standard emergency room screening evaluations alone or to standard evaluations plus cardiac CT angiography.

The researchers studied 1,000 participants between 40 and 74 years old in nine U.S. hospitals. Participants were eligible to enroll in the trial if they showed symptoms suggestive of acute coronary syndrome but no prior history of heart disease or evidence of heart damage on their electrocardiogram (ECG) tests or blood tests.

"The results from this study should help health care providers and patients make better informed decisions by knowing the risks and potential benefits of using CT scans to more quickly diagnose [acute coronary syndrome](#)," said Udo Hoffmann, M.D., M.P.H., the study's principal investigator at Massachusetts General Hospital and Harvard Medical School, Boston. "It can be a relief to [patients](#) with chest pain to quickly know they are not having a [heart](#) attack and that they can spend the night at home, instead of in a hospital bed."

Participants in the CT group were exposed to more radiation than those in the standard screening group, though the study authors suggested that future CT scans could be done using less radiation, which could help lower exposure without sacrificing accuracy.

Overall costs were similar in the two treatment groups.

**More information:** Learn more about this clinical trial (NCT01084239) at: [clinicaltrials.gov/ct2/show/NCT01084239](https://clinicaltrials.gov/ct2/show/NCT01084239)

Provided by NIH/National Heart, Lung and Blood Institute

Citation: Heart CT scans may help emergency room personnel more quickly assess patients with chest pain (2012, July 25) retrieved 11 July 2024 from <https://medicalxpress.com/news/2012-07-heart-ct-scans-emergency-room.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.