

HPTN study finds greatly elevated HIV infection rates among young black MSM in the US

July 23 2012

Study results released today by the HIV Prevention Trials Network (HPTN) show disturbing rates of new HIV infections occurring among black gay and bisexual men in the U.S. (also known as men who have sex with men, or MSM), particularly young black MSM. The HPTN 061 study showed that the overall rate of new HIV infection among black MSM in this study was 2.8% per year, a rate that is nearly 50% higher than in white MSM in the U.S. Even more alarming, HPTN 061 found that young black MSM—those 30 years of age and younger—acquired HIV infection at a rate of 5.9% per year, three times the rate among U.S. white MSM. The overall infection rate among black MSM in this U.S. study is comparable to the rate seen in the general populations of countries in sub-Saharan Africa hardest hit by the HIV epidemic. HPTN 061 was a large multi-site study of HIV and black MSM conducted in six U.S. cities, and the first to determine the rate of new HIV infection among such a large prospective cohort of U.S. black MSM (referred to as HIV incidence).

Commenting on the findings, HPTN 061 Study Co-Chair Darrell Wheeler, PhD, MPH, Dean of the School of Social Work at Loyola University said, "We have known that black MSM are affected by [HIV](#) at disproportionately higher rates when compared to other MSM in the U.S., but the HPTN 061 HIV incidence rates were extremely high. They make it very clear that we must urgently find and implement ways to stem the spread of HIV among [black gay](#) men in this country, and

critically among young black gay men."

The HIV incidence results from HPTN 061, presented today at the XIX International AIDS Conference in Washington, D.C. are among the first findings to come out of the study. Other early analyses show that [HIV infection](#) in this study population was associated with very high rates of untreated sexually transmitted infections and with poverty. Of the self-reported HIV-negative or unknown HIV status study participants who agreed to be tested at time of enrollment, 12% were HIV-positive. This finding suggests low rates of HIV testing or delays in testing among black MSM. All HIV-infected study participants were referred for HIV care and treatment services. During the study, all participants were offered the chance to work with "peer health navigators" to provide support and to assist them with access to needed services. In addition, they were provided with condoms, testing and referral treatment of sexually transmitted infections, risk reduction counseling and other standard HIV prevention services.

The success of the HPTN 061 in recruiting and retaining black MSM in a year-long study, including 97% uptake of HIV testing, answered a primary objective of the study, and paves the way for the conduct of future HIV prevention research.

"The HPTN 061 study findings are a sobering wake-up call," said Wafaa El-Sadr, MD, MPH Co-Principal Investigator for the HPTN. "These results, and others to be garnered from further analysis of the study, highlight the enormity of the challenges ahead and offer important insights into how to design research studies and programs for prevention of HIV among black MSM."

Further analysis of data from the study is ongoing to assess how HIV risk among [black](#) MSM might be affected by childhood experiences, social and sexual networks, discrimination, homophobia, incarceration and

barriers to health care. These findings will help inform future research and prevention efforts.

The HPTN 061 study enrolled a total of 1,553 MSM in six U.S. cities; Atlanta, GA, Boston, MA, New York, NY, Los Angeles, CA, San Francisco, CA, and Washington, D.C. and was conducted between 2009-2011. The study is funded by the National Institute of Allergy and Infectious Diseases (NIAID), the National Institute on Drug Abuse (NIDA) and the [National Institute](#) of Mental Health (NIMH).

Provided by HIV Prevention Trials Network (HPTN)

Citation: HPTN study finds greatly elevated HIV infection rates among young black MSM in the US (2012, July 23) retrieved 26 April 2024 from <https://medicalxpress.com/news/2012-07-hptn-greatly-elevated-hiv-infection.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.