

Cholesterol levels appear to be improving among US youths

August 7 2012

In a study involving more than 16,000 U.S. children and adolescents, there has been a decrease in average total cholesterol levels over the past 2 decades, although almost 1 in 10 had elevated total cholesterol in 2007-2010, according to a study in the August 8 issue of *JAMA*.

"The process of atherosclerosis begins during childhood and is associated with adverse serum lipid concentrations including high concentrations of low-density lipoprotein cholesterol (LDL-C), non-highdensity lipoprotein cholesterol (non-HDL-C), and triglycerides, and low concentrations of high-density lipoprotein cholesterol (HDL-C). Serum lipid concentrations in childhood are associated with serum lipid concentrations in adulthood," according to background information in the article. "For more than 20 years, primary prevention of coronary heart disease has included strategies intended to improve overall serum lipid concentrations among youths."

Brian K. Kit, M.D., M.P.H., of the <u>Centers for Disease Control and</u> <u>Prevention</u>, Hyattsville, Md., and colleagues conducted a study to examine the trends in serum lipid concentrations among children and adolescents in the U.S. The study included data on 16,116 youths ages 6 to 19 years who participated in the nationally representative National Health and <u>Nutrition Examination Survey</u> (NHANES) during 3 time periods: 1988-1994,1999-2002, and 2007-2010. Among the measures analyzed for all the participants were average serum total cholesterol (TC), non-HDL-C, HDL-C; and among adolescents only (ages 12-19 years), LDL-C and geometric average <u>triglyceride levels</u>. Trends in



adverse lipid concentrations were reported for TC levels of 200 mg/dL and greater, non-HDL-C levels of 145 mg/dL and greater, HDL-C levels of less than 40 mg/dL, LDL-C levels of 130 mg/dL and greater, and triglyceride levels of 130 mg/dL and greater.

The researchers found that among youths ages 6 to 19 years between 1988-1994 and 2007-2010, there was a decrease in average serum TC from 165 mg/dL to 160 mg/dL and an increase in average serum HDL-C from 50.5 mg/dL to 52.2 mg/dL. There was also a decrease in the average serum non-HDL-C levels during this time period. "Generally, the sex-, age-, and race/ethnicity-specific trends for TC, HDL-C, and non-HDL-C were similar in direction to the overall trends and consistent with a favorable trend, although for each group, the magnitude was not the same and the trend was not always significant," the authors write.

Between 1988-1994 and 2007-2010, there was a decrease in prevalence among youths ages 6 to 19 years of elevated TC from 11.3 percent to 8.1 percent and non-HDL-C from 13.6 percent to 10 percent. Prevalence of low HDL-C was 17.3 percent in 1988-1994 and 14.8 percent in 2007-2010, a nonsignificant decrease. Among adolescents, there was a decrease in prevalence of elevated LDL-C and triglycerides between 1988-1994 and 2007-2010 and a decrease in average serum LDL-C from 95 mg/dL to 90 mg/dL and geometric average serum triglycerides from 82 mg/dL to 73 mg/dL.

In 2007-2010, 22 percent of youths had either a low HDL-C level or high non-HDL-C, which was lower than the 27.2 percent in 1988-1994.

The authors also found that age- and race/ethnicity-adjusted TC was 4.3 mg/dL lower for males and 6.5 mg/dL lower for females in 2007-2010 than in 1988-1994. Males and females who were non-Hispanic black or Mexican American, and also females who were non-Hispanic white had a lower age-adjusted TC in 2007-2010 than in 1988-1994. Age- and



race/ethnicity- adjusted HDL-C was higher for males and females in 2007-2010 than in 1988-1994.

"Between 1988-1994 and 2007-2010, a favorable trend in serum lipid concentrations was observed among youths in the United States but adverse lipid profiles continue to be observed among youths," the researchers write. "For example, in 2007-2010, slightly more than 20 percent of children aged 9 to 11 years had either a low HDL-C or high non-HDL-C concentration, which, according to the most recent cardiovascular health guidelines for children and adolescents, indicates a need for additional clinical evaluation."

"The recently released Expert Panel on Integrated Guidelines for Cardiovascular Health and Risk Reduction in Children and Adolescents provides recommendations for preventing the development of cardiovascular risk factors including optimizing nutrition and physical activity and reducing exposure to tobacco smoke. Specific screening approaches, including universal screening at select ages and management of adverse lipid concentrations, have also been detailed in these recent guidelines for youths. Future research from longitudinal studies or mortality-linked data, including NHANES, may include examining clinical outcomes for cardiovascular disease, including cardiovascular mortality, based on lipid concentrations present during childhood," the authors conclude.

Sarah D. de Ferranti, M.D., M.P.H., of Harvard Medical School and Boston Children's Hospital, Boston, comments on the findings of this study in an accompanying editorial.

"Further research is needed to investigate the hypothesized contributors to change in childhood and adolescent lipid levels. Better understanding is needed about dietary trends and physical activity during childhood, areas that were not explored in the study by Kit et al but could be



assessed using NHANES data. However it seems clear that populationwide efforts to alter cardiovascular disease [CVD] risk have potential to influence health risks. Examples of other interventions that could have positive population-level effects on health include taxation of carbonated beverages, improved access to water in schools and in the workplace, and environmental changes to promote daily physical activity. Improvements in child and adolescent lipid values over the past 2 decades are significant and may portend improved CVD outcomes for the future, but much work should be done to better understand the changes and to build upon them."

More information:

JAMA. 2012;308[6]:591-600. *JAMA*. 2012;308[6]:621-622.

Provided by JAMA and Archives Journals

Citation: Cholesterol levels appear to be improving among US youths (2012, August 7) retrieved 7 May 2024 from <u>https://medicalxpress.com/news/2012-08-cholesterol-youths.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.