

## In epileptic seizures, researchers see the neurology of consciousness

August 15 2012, By Bill Hathaway



The fMRI images are different viewpoints of the brain of a child experiencing an epileptic seizure. Areas in yellow and orange represent increased brain activity compared to its normal state, while areas in blue show decreased activity. These are the areas of the brain needed for normal consciousness.

(Medical Xpress) -- Yale researchers studying epileptic seizures have shed new light on the neurological origins of consciousness.



When epileptics lose consciousness during a variety of types of seizures, the signals converge on the same brain structures, but through different pathways, says Dr. Hal Blumenfeld, professor of neurology, <u>neurobiology</u>, and neurosurgery.

"Understanding of these mechanisms could lead to improved treatment strategies to prevent impairment of consciousness and improve the quality of life of people with epilepsy," he said.

Blumenfeld's research is described in the current issue of the journal *Lancet Neurology*.

Provided by Yale University

Citation: In epileptic seizures, researchers see the neurology of consciousness (2012, August 15) retrieved 5 May 2024 from https://medicalxpress.com/news/2012-08-epileptic-seizures-neurology-consciousness.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.