

# 'Harmless' condition shown to alter brain function in elderly

August 13 2012

---

Researchers at the Mayo Clinic say a common condition called leukoaraiosis, made up of tiny areas in the brain that have been deprived of oxygen and appear as bright white dots on MRI scans, is not a harmless part of the aging process, but rather a disease that alters brain function in the elderly. Results of their study are published online in the journal *Radiology*.

"There has been a lot of controversy over these commonly identified abnormalities on MRI scans and their clinical impact," said Kirk M. Welker, M.D., assistant professor of radiology in the College of Medicine at Mayo Clinic in Rochester, Minn. "In the past, leukoaraiosis has been considered a benign part of the aging process, like gray hair and wrinkles."

Leukoaraiosis, also called small vessel ischemia and often referred to as unidentified bright objects or "UBOs" on [brain scans](#), is a condition in which diseased blood vessels lead to small areas of damage in the white matter of the brain. The lesions are common in the brains of people over the age of 60, although the amount of disease varies among individuals.

"We know that aging is a risk factor for leukoaraiosis, and we suspect that high blood pressure may also play a role," Dr. Welker said.

Dr. Welker's team performed functional MRI (fMRI) scans on cognitively normal elderly participants recruited from the Mayo Clinic Study of Aging between 2006 and 2010. In 18 participants, the amount

of leukoaraiosis was a moderate 25 milliliters, and in 18 age-matched [control participants](#), the amount of disease was less than five milliliters.

The patients were imaged in an [MRI scanner](#) as they performed a semantic decision task by identifying [word pairs](#) and a [visual perception](#) task that involved differentiating straight from diagonal lines. fMRI is a special type of [magnetic resonance imaging](#) that measures metabolic changes in an active part of the brain.

Although both groups performed the tasks with similar success, the fMRI scans revealed different [brain activation](#) patterns between the two groups. Compared to members of the control group, patients with moderate levels of leukoaraiosis had atypical activation patterns, including decreased activation in areas of the brain involved in language processing during the semantic decision task and increased activation in the visual-spatial areas of the brain during the visual perception task.

"Different systems of the brain respond differently to disease," Dr. Welker explained. "White matter damage affects connections within the brain's language network, which leads to an overall reduction in network activity."

He pointed out that identifying leukoaraiosis in the brain is important, both for individual patients undergoing brain mapping for surgery or other treatments and for research studies.

For improved neurological health, Dr. Welker said efforts should be taken to prevent leukoaraiosis from occurring.

"Our results add to a growing body of evidence that this is a disease we need to pay attention to," he said. "Leukoaraiosis is not a benign manifestation of aging but an important pathologic condition that alters [brain function](#)."

**More information:** "Altered Functional MR Imaging Language Activation in Elderly Individuals with Cerebral Leukoaraiosis," *Radiology*.

Provided by Radiological Society of North America

Citation: 'Harmless' condition shown to alter brain function in elderly (2012, August 13)  
retrieved 3 May 2024 from

<https://medicalxpress.com/news/2012-08-harmless-condition-shown-brain-function.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--