

Intervention helps children with sickle cell disease complete MRI tests without sedation

August 31 2012



St. Jude Children's Research Hospital reports success in using a targeted educational approach to teach young sickle cell disease patients to remain motionless during MRI scans, making the process safer. Credit: St. Jude Biomedical Communications

Sitting still is tough for children, which makes MRI scans a challenge. The scans require that patients remain motionless for extended periods. Findings from St. Jude Children's Research Hospital showed that a brief, targeted intervention dramatically increases the likelihood that children as young as 5 years old will be able to undergo testing without sedation.

That is good news for children with sickle cell disease, who were the focus of this study. [Patients](#) with sickle cell disease often undergo brain and liver MRIs to check for complications related to their disease or treatment. But the [blood disorder](#) puts patients at increased risk for

serious anesthesia-related complications, so [sedation](#) is avoided when possible.

This study of 71 patients with sickle cell disease ages 5 through 12 found that children who completed the short preparation program prior to their MRI were eight times more likely to complete the scans without being sedated than patients of the same age who did not receive the preparation. The work was published online in the journal *Pediatric Radiology*.

The intervention implemented by the Child Life Program at St. Jude involves educating patients and families about exactly what to expect during an MRI, assigning patients jobs to focus on during the scans and also working to identify strategies to help them remain motionless during the test. "Some patients chose to listen to music or to squeeze a ball to help them remember not to move," said Katherine Cejda, a St. Jude Child Life specialist and the study's first author. "Some patients had the option of watching movies or having parents or other [adults](#) in the room with them during the test."

This study is the first designed to determine the effectiveness of this approach and the first to focus on children with sickle cell disease. Cejda said similar programs are used by child life specialists throughout the U.S. to help prepare children for MRIs and other [medical procedures](#). September is National Sickle Cell Awareness Month. This study highlights the varied efforts underway at St. Jude to improve the health and quality of life for children and adolescents with sickle cell disease.

The paper's senior author, Jane Hankins, M.D., an associate member of the St. Jude Department of Hematology, said that avoiding sedation reduces the risk to the patients, particularly those with sickle cell disease.

Sickle cell disease is caused by a mutation in the gene for assembling

hemoglobin. Hemoglobin is the protein red blood cells use to carry oxygen throughout the body. While red blood cells are normally pliable discs, the inherited mistake results in red blood cells becoming brittle and assuming a banana shape that can trigger intense pain and lead to organ damage. Hankins said the disease also makes anesthesia riskier.

General anesthesia can lead to a drop in temperature, blood oxygen levels and dehydration. In patients with sickle cell disease, dehydration can result in a painful and dangerous buildup of sickled cells. Patients can wind up hospitalized, sometimes in the intensive care unit, for treatment of severe pain or acute chest syndrome, a pneumonia-like illness, Hankins said. To reduce that risk, a sickle cell patient scheduled for anesthesia is hospitalized overnight to receive extra fluids and possibly a blood transfusion.

The combination of factors creates inconvenience and added risk for the patient. It also adds to health care costs. While no family ever pays St. Jude for the care their child receives, controlling costs remains important to the hospital. The program could also be adopted by other health care institutions worldwide to help manage cost.

The intervention is now routinely offered to all St. Jude patients undergoing MRIs. Cejda said the program has been used to help children as young as age 4 complete scans without sedation. "This preparation program offers a real advantage to patients," Hankins added.

The study included children scheduled for [brain](#) or [liver](#) MRIs, which usually last 30 to 60 minutes. Researchers reported that 30 of the 33 [children](#), or 91 percent, in the intervention group underwent successful MRI scans without sedation. That compares to 71 percent, or 27 of 33 patients, who did not participate in the program prior to undergoing MRIs without sedation. The other authors are Matthew Smeltzer, Beth McCarville and Kathleen Helton, all of St. Jude, and Eileen Hansbury,

formerly of St. Jude.

Provided by St. Jude Children's Research Hospital

Citation: Intervention helps children with sickle cell disease complete MRI tests without sedation (2012, August 31) retrieved 25 April 2024 from

<https://medicalxpress.com/news/2012-08-intervention-children-sickle-cell-disease.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.