

New study: Running mechanics, not metabolism, are the key to performance for elite sprinters

August 1 2012



Sprinters competing in the 2012 Olympics might assume their championship performance is the result of their fuel-efficient physiology.

But a new study disproves the classic scientific view that [conserving energy](#) maximizes [performance](#) in a sprinting event.

The study by biomechanics researchers Matthew W. Bundle at the University of Montana and Peter G. Weyand at Southern Methodist University, Dallas, demonstrates that metabolic economy is not an

important factor for performance in events lasting 60 seconds or less.

In fact, just the opposite is true.

"That prevailing view is no longer viable," said Weyand. "Sprinters, if anything, are wasteful of energy. This is due to the biological trade-offs between faster muscle fibers that provide the large and rapid forces needed for sprinting, and slower muscle fibers that maximize metabolic economy."

Instead, the key to top-flight sprinting is to maximize how hard each foot hits the ground, which allows sprinters to translate musculoskeletal and ground reaction forces into swift motion, said Bundle.

"[Saving energy](#) is critically important for endurance, but not for sprinting, which our findings indicate is not energy-limited," Bundle said.

[Metabolic energy](#) available from sustainable, aerobic sources predominantly determines performance during endurance events by setting the intensity of the musculoskeletal performance that can be sustained throughout the effort, the study found.

For sprinters, Bundle and Weyand conclude the opposite is true.

"The intensity of the mechanical activity that the musculoskeletal system can (for a very short time) achieve determines the quantities of metabolic energy released and the level of performance attained," according to the study.

The authors reported their findings in "Sprint [Exercise Performance: Does Metabolic Power Matter?](#)" in the July issue of *Exercise and [Sport Sciences Reviews](#)*, <http://bit.ly/ODvCrk>.

Sprint performance variations are a function of external forces

The authors write in their study that athletic performance can be analyzed considering either the input to, or the output from, the skeletal muscles that serve as biological engines. Input is the chemical energy that fuels muscular contraction. Output is the force or mechanical power the contractions produce.

To analyze the mechanics of burst-type sprint activities, the authors said they drew on all-out running speeds and cycling power outputs of humans because of the abundance and quality of the data available and because the mechanical and metabolic contrasts between the two provide informative insights. The authors focused on durations of up to five minutes, particularly on efforts of less than a minute.

For both exercises, differences in sprinting performance were predominantly a function of the magnitude of the external forces applied because running contact lengths and cycling down-stroke lengths, as well as stride and pedal frequency, exhibited limited variations. Additionally, for both cycling and running, external forces applied during sprinting are believed to be consistently related to the corresponding muscle forces, regardless of the intensity or duration of the effort.

So what determines the maximum external forces the [musculoskeletal system](#) can apply during a brief, all-out sprint? And why do those forces decrease over the duration of the sprint?

The researchers assessed neuromuscular activation using a diagnostic procedure called surface electromyography to measure electrical activity in the activated [muscle fibers](#). That assessment showed that neuromuscular activation increases continuously during all-out sprint

cycling and running trials. More rapid increases were typical for the briefest trials that required the greatest forces. That indicates that all-out sprinting performances are highly dependent on duration because of the speed of musculoskeletal fatigue during dynamic exercise requiring large force outputs, the authors reported.

Sprint performance linked to mechanics of applying external force

Bundle and Weyand altered three independent variables to maximize the variation observed in sprint performance: Subjects were individuals with large differences in their sprint performance capabilities; all-out sprint trials spanned a broad range of durations from 2 to 300 seconds; and performance was compared across different modes of sprinting, namely cycling and running.

"The predictive success of our force application model, both within and across modes of sprint exercise, indicates that as efforts extend from a few seconds to a few minutes, the fractional reliance on anaerobic metabolism progressively impairs whole-body musculoskeletal performance, and does so with a rapid and remarkably consistent time course," the authors wrote. "In this respect, the sprint portion of the performance-duration curve predominantly represents, not a limit on the rates of energy re-supply, but the progressive impairment of skeletal muscle force production that results from a reliance on anaerobic metabolism to fuel intense, sequential contractions."

Conclusion of study departs from prevailing physiological paradigm

Since the muscular engines of humans and other animals are similar in terms of their metabolic and mechanical function, the findings likely

apply to the burst performance capabilities of vertebrate animals in general, say the researchers.

More information: journals.lww.com/acsm-essr/Abs...abolic_Power.10.aspx

Provided by Southern Methodist University

Citation: New study: Running mechanics, not metabolism, are the key to performance for elite sprinters (2012, August 1) retrieved 27 April 2024 from <https://medicalxpress.com/news/2012-08-mechanics-metabolism-key-elite-sprinters.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.