

Even mild depression, anxiety hurts the heart: study

August 1 2012, By Steven Reinberg, HealthDay Reporter



The risk rises as chronic mental distress becomes more intense, study says.

(HealthDay) -- Even mild depression or anxiety may raise your risk of dying from cardiovascular disease and other causes, according to British researchers.

And the greater the level of psychological distress, the higher the odds of death from [heart disease](#), the researchers say.

"The fact that an increased risk of mortality was evident, even at low levels of psychological distress, should prompt research into whether treatment of these very common, minor symptoms can reduce this increased risk of death," said lead researcher Tom Russ, a clinical research fellow at the Alzheimer Scotland [Dementia](#) Research Center of

the University of Edinburgh.

For the study, published online July 31 in [BMJ](#), Russ and colleagues analyzed 10 studies of men and women enrolled in the [Health Survey](#) for England from 1994 to 2004. Data on more than 68,000 adults aged 35 and older was included overall.

Each study looked for connections between chronic psychological distress and the risk of dying from heart disease and other causes, including cancer.

Pooling data in this way is called a meta-analysis. In such a study, researchers look for common patterns across several studies.

Over eight years' follow-up, the researchers found even very [mild depression](#) or [anxiety](#) -- subclinical levels -- raised the risk of all-cause death, including [cardiovascular disease](#), by 20 percent. Looking specifically at death from heart disease, mild psychological distress raised this risk 29 percent, the study found.

For the highest level of [depression](#) or anxiety, the risk of all-cause death rose 94 percent, the researchers found.

Risk of death from cancer was increased 9 percent in cases of very [severe depression](#) or anxiety, the investigators found. Lower levels of psychological distress were not associated with increased risk of [cancer death](#).

An individual's actual risk of death remains small, however, and people shouldn't assume they are doomed to an early [death](#) if they suffer from a psychological disorder.

Dr. Glyn Lewis, a professor of psychiatric epidemiology at the

University of Bristol in England and author of an accompanying journal editorial, said evidence linking stress to heart disease continues to mount.

"If we can reduce the psychological impact, then this should reduce the biological response," he said. But how to accomplish that remains a puzzle.

A type of psychological treatment called cognitive behavioral therapy is designed to help people change the way they respond to potentially stressful events, Lewis said. Cognitive behavioral therapy teaches patients to change their thinking about situations and to react less emotionally.

"This might help people with [clinical] depression, but there is no evidence that this might help the much larger numbers of people who have low-level symptoms that are below the diagnostic threshold for depression," he said.

While antidepressants might improve depression, previous studies have linked their use to greater risk of heart disease, according to background research in the study. About 7.5 percent of United Kingdom residents have depression and anxiety disorders, Lewis said.

Changing this stress-disease dynamic might also involve keeping common risk factors for cardiovascular disease in check, another expert said.

Dr. Gregg Fonarow, a professor of cardiovascular medicine at the University of California, Los Angeles, said many studies have demonstrated an association between depression and anxiety and cardiovascular events, cardiovascular deaths and all-cause mortality.

But so far, no evidence has shown that treating depression or anxiety reduces the risk of heart disease, Fonarow said.

Many different mechanisms may connect [psychological distress](#) to cardiovascular disease, including increased sympathetic nervous system activity, stress hormones such as cortisol, chronic inflammation, unhealthy lifestyle factors and inattention to early symptoms, he said.

"For people with depression or anxiety, focusing on proven cardiovascular risk factor interventions, including maintaining healthy blood pressure, body weight, cholesterol levels, engaging in regular exercise and not smoking, may represent the best course of action to lower their cardiovascular risk," he advised.

More information: For more information on cardiovascular disease, visit the [American Heart Association](#).

Copyright © 2012 [HealthDay](#). All rights reserved.

Citation: Even mild depression, anxiety hurts the heart: study (2012, August 1) retrieved 18 April 2024 from <https://medicalxpress.com/news/2012-08-mild-depression-anxiety-heart.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--