

Renal denervation gives better outcomes than drugs in advanced heart failure

August 27 2012

Renal denervation leads to better outcomes than standard drug treatment in patients with advanced heart failure, according to research presented at ESC Congress 2012. The results of the Olomouc I pilot study were presented by Dr Miloš Táborský from the University Hospital Olomouc.

Renal [denervation](#) does not involve any pharmacological treatment. It is a multiple application of radiofrequency energy using a thin [catheter](#) via the [femoral artery](#) and a long-term "denervation" of the [sympathetic nerves](#) around the renal arteries. "This is done to prevent an increased [excretion](#) of the substances leading to heart failure," said Dr Táborský.

The aim of this pilot study was to compare the outcomes of patients with advanced heart failure receiving standard pharmacological treatment or renal denervation. Over the period of one year, 26 patients were treated with renal denervation and standard [pharmacotherapy](#) and 25 patients received standard drug treatment with beta-blockers, [ACE inhibitors](#) or ARBs and diuretics.

After one year contractile function of the left side of the heart improved in patients treated with renal denervation from $25\pm 12\%$ to $31\pm 14\%$ (p

Citation: Renal denervation gives better outcomes than drugs in advanced heart failure (2012, August 27) retrieved 20 April 2024 from <https://medicalxpress.com/news/2012-08-renal-denervation-outcomes-drugs-advanced.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.