

Study reveals optimal treatment for most common infection after organ transplantation

August 23 2012

Waiting to treat the commonest viral infections in transplant recipients until they reach a certain threshold is better than prophylactically treating all recipients, according to a study appearing in an upcoming issue of the *Journal of the American Society of Nephrology* (JASN).

Cytomegalovirus (CMV) infection is the most common infection in [organ transplant recipients](#), who are susceptible to infections in general because they must take immunosuppressive medications long term. CMV infections can cause increased risks of other infections, [organ rejection](#), [heart complications](#), and diabetes.

The two main strategies against CMV are called universal antiviral prophylaxis and pre-emptive therapy. In universal prophylaxis all patients at risk are given [antiviral drugs](#) (such as [valganciclovir](#), valacyclovir, and ganciclovir) for several months after transplantation. In pre-emptive therapy, patients are intensively monitored for CMV activity by sensitive laboratory methods, and short-term [antiviral treatment](#) is given only to those with significant viral counts before symptoms occur.

To compare these strategies, Tomas Reischig, MD, PhD (Charles University Medical School and Teaching Hospital, in Pilsen, Czech Republic) and his colleagues assigned [kidney transplant recipients](#)—who either had CMV present in their blood or who received transplants from

donors with CMV in their blood—to either three months of prophylaxis with valacyclovir or pre-emptive valganciclovir given when significant CMV counts were detected.

Among the major findings for 55 patients after three years:

- CMV disease developed in 6% of patients in the pre-emptive therapy group and in 9% receiving prophylaxis
- Patients in the prophylaxis group were 2.5 times more likely to develop moderate-to-severe kidney scarring and atrophy than patients receiving pre-emptive therapy.
- Kidney biopsies showed that the prophylaxis group also had significantly higher expression of genes involved in kidney scarring.
- The occurrence of CMV was similar in both groups, but pre-emptive therapy improved 4-year survival of transplanted organs (92% vs 74%).

These findings indicate that compared with valacyclovir prophylaxis, pre-emptive valganciclovir therapy for CMV may lead to less severe kidney scarring and atrophy and to significantly better survival of transplanted organs.

"In the view of short-term trial results, which favor CMV prophylaxis over pre-emptive strategy because of lower risk of acute rejection, we expected a translation of presumed benefit of prophylaxis to the long-term post-transplant period. In fact, we discovered that the opposite is true," said Dr. Reischig.

More information: [doi: 10.1681/ASN.2012010100](https://doi.org/10.1681/ASN.2012010100)

Provided by American Society of Nephrology

Citation: Study reveals optimal treatment for most common infection after organ transplantation (2012, August 23) retrieved 27 April 2024 from

<https://medicalxpress.com/news/2012-08-reveals-optimal-treatment-common-infection.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.