

Fasting makes brain tumors more vulnerable to radiation therapy

September 11 2012

A new study from USC researchers is the first to show that controlled fasting improves the effectiveness of radiation therapy in cancer treatments, extending life expectancy in mice with aggressive brain tumors.

Prior work by USC professor of gerontology and biological sciences Valter Longo, corresponding author on the study and director of the Longevity Institute at the USC Davis School of [Gerontology](#), has shown that short-term fasting protects healthy cells while leaving [cancer cells](#) vulnerable to the toxic effects of chemotherapy.

The latest study, appearing online once the embargo lifts in *PLOS One*, is the first to show that periods of fasting appear to have the same augmenting effect on [radiation therapy](#) in treating gliomas, the most commonly diagnosed brain tumor. Gliomas have a median survival of less than two years.

"With our initial research on chemotherapy, we looked at how to protect patients against toxicity. With this research on radiation, we're asking, what are the conditions that make cancer most susceptible to treatment? How can we replicate the conditions that are least hospitable to cancer?" Longo said.

Longo and his co-investigators studied the combination of fasting with radiation therapy and with the chemotherapy drug Temozolomide, currently the standard treatment for the treatment of brain tumors in

adults after an attempt at surgical removal.

They found that controlled short-term fasting in mice, no more than 48 hours each cycle, improved the effectiveness of radiation and chemotherapy in treating gliomas. Despite the extremely aggressive growth of the type of brain tumor studied, more than twice as many mice who fasted and received radiation therapy survived to the end of the trial period than survived with radiation alone or fasting alone.

"The results demonstrate the beneficial role of fasting in gliomas and their treatment with standard [chemotherapy](#) and radiotherapy," the researchers write, who say the results indicate the benefits of short-term, controlled fasting for humans receiving treatment for [brain tumors](#).

Longo cautions that patients should consult with their oncologist before undertaking any fasting: "You want to balance the risks. You have to do it right. But if the conditions are such that you've run out of options, short-term fasting may represent an important possibility for patients."

Provided by University of Southern California

Citation: Fasting makes brain tumors more vulnerable to radiation therapy (2012, September 11) retrieved 24 April 2024 from

<https://medicalxpress.com/news/2012-09-fasting-brain-tumors-vulnerable-therapy.html>

| |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|