

Study illuminates roles of novel epigenetic chemical in the brain

September 17 2012

Researchers from the Centre for Addiction and Mental Health (CAMH) have identified a new role of a chemical involved in controlling the genes underlying memory and learning.

"The brain is a [plastic](#) tissue, and we know that [learning and memory](#) require various genes to be expressed," says CAMH Senior Scientist Dr. Art Petronis, who is a senior author on the new study. "Our research has identified how the chemical 5-hmC may be involved in the epigenetic processes allowing this plasticity." Dr. Petronis is head of the Krembil Family Epigenetics Laboratory in CAMH's Campbell Family Mental Health Research Institute.

5-hmC is an epigenetic modification of DNA, and was discovered in humans and mice in 2009. DNA modifications are [chemical changes](#) to DNA. They flag genes to be turned "on" – signalling the genome to make a protein – or turned "off." As the overwhelming majority of cells in an individual contain the same [genetic code](#), this pattern of flags is what allows a neuron to use the same genome as a blood or [liver cell](#), but create a completely different and specialized [cellular environment](#).

The research, published online in *Nature Structural & Molecular Biology*, sheds light on the role of 5-hmC. Intriguingly, it is more abundant in the brain than in other tissues in the body, for reasons not clear to date.

The CAMH team of scientists examined DNA from a variety of tissues, including the mouse and human brain, and looked at where 5-hmC was

found in the genome. They detected that 5-hmC had a unique distribution in the brain: it was highly enriched in genes related to the synapse, the dynamic tips of brain cells. Growth and change in the synapse allow different brain cells to "wire" together, which enables learning and memory.

"This enrichment of 5-hmC in synapse-related genes suggests a role for this epigenetic modification in learning and memory," says Dr. Petronis.

The team further showed that 5-hmC had a special distribution even within the gene. The code for one gene can be edited and "spliced" to create several different proteins. Dr. Petronis found that 5-hmC is located at "splice junctions," the points where the gene is cut before splicing.

"5-hmC may signal the cell's splicing machinery to generate the diverse proteins that, in turn, give rise to the unprecedented complexity of the brain," he says.

The research team is continuing to investigate the role of 5-hmC in more detail, and to determine whether 5-hmC function is different in people with bipolar disorder and schizophrenia compared to people without these diagnoses.

Provided by Centre for Addiction and Mental Health

Citation: Study illuminates roles of novel epigenetic chemical in the brain (2012, September 17) retrieved 5 May 2024 from

<https://medicalxpress.com/news/2012-09-illuminates-roles-epigenetic-chemical-brain.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
