

# WHO urges action as drug-resistant malaria spreads

September 27 2012

---

The World Health Organization said Thursday that governments in the Mekong region must act "urgently" to stop the spread of drug-resistant malaria which has emerged in parts of Vietnam and Myanmar.

There is growing evidence that the [malaria parasite](#) is becoming resistant to a frontline treatment, the anti-malarial drug artemisinin, in southern and central Vietnam and in southeastern Myanmar, the WHO said in a statement.

Drug-[resistant malaria](#) was found on the Thailand-Cambodia border eight years ago, and has since also been discovered along the Thailand-Myanmar border, according to scientists.

WHO regional director, Shin Young-soo, said countries must "urgently address this issue before we put at risk not only the fragile gains we have made in [malaria control](#) but also our goal of a malaria-free Western Pacific Region."

Countries in the Mekong region must "intensify and expand" operations to contain and eliminate artemisinin-resistant malaria, Shin said at a WHO regional meeting in Hanoi.

Drug-resistant malaria in Vietnam is mainly concentrated in three central and southern provinces, say health officials, adding that there are a number of key factors which make tackling the problem difficult in the communist country.

The population in the affected provinces are very poor, highly mobile, and have not widely embraced DEET-treated [mosquito nets](#) which could reduce malaria infections, according to media reports of a government conference in July.

People in the affected provinces often received outpatient treatment and frequently fail to complete a full course of anti-malarial treatment.

Resistance to [artemisinin](#) does not prevent patients being cured, thanks to other partner drugs, but treatment typically takes a few days instead of 24 hours.

Malaria is a life-threatening disease caused by parasites that are transmitted through the bites of infected mosquitoes.

It killed an estimated 655,000 people in 2010, mostly African children, although mortality rates have fallen sharply over the past decades, according to the WHO.

(c) 2012 AFP

Citation: WHO urges action as drug-resistant malaria spreads (2012, September 27) retrieved 18 April 2024 from

<https://medicalxpress.com/news/2012-09-urges-action-drug-resistant-malaria.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--