

Children with obesity must be treated in time, prior to adolescence, researchers warn

October 30 2012

A new study from Karolinska Institutet shows that behavioural treatment for inducing weight-loss can be very effective for severely obese children. However, the treatment to change dietary and exercise habits must be given in time, as it showed to have little effect on adolescents with the same problem.

"The results are indeed alarming for these severely obese adolescents, who run a serious risk of disease and social marginalisation," says Claude Marcus, professor of paediatrics at Karolinska Institutet. "New treatment methods must be developed and for some young people, surgery must even be considered an option."

In the present study, which is published in the scientific journal *Archives of Pediatrics & Adolescent Medicine*, the researchers have evaluated a form of behavioural treatment for seriously obese children centred on exercise and diet. Over a period of three years, the children met regularly with a team comprising a doctor, psychologist, physiotherapist, nurse and dietician at Rikshälsan Barnobesitas, a national childhood obesity centre based at the Astrid Lindgren Childrens Hospital in Stockholm. The study included 643 obese children and adolescents in the 6-16 age-band, who began treatment between 1998 and 2006.

The results show that the behavioural treatment was effective for the younger children, particularly those with severe obesity. But the older the participants were, the harder it became to change their dietary and [exercise habits](#). Teenagers with moderate obesity responded slightly to

the treatment, but for the large majority of teenagers with severe obesity, it had no demonstrable health effect at all.

The study also revealed that most of the obese [adolescents](#) had weight problems already in primary school, leading the researchers to conclude that much would be gained if treatment was started when overweight or obese children were six or seven years old.

"Unfortunately, data from the Swedish national Childhood [Obesity](#) Registry, BORIS, show that the average age for beginning treatment is currently ten," says study team member Dr Pernilla Danielsson. "And there are some health authorities that offer no treatment at all to [obese children](#)."

More information: Danielsson, P. et al., Response of Severely Obese Children and Adolescents to Behavioral Treatment. *Archives of Pediatrics & Adolescent Medicine*, first online 29 October. 2012

Provided by Karolinska Institutet

Citation: Children with obesity must be treated in time, prior to adolescence, researchers warn (2012, October 30) retrieved 9 April 2024 from <https://medicalxpress.com/news/2012-10-children-obesity-prior-adolescence.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--