

A plan to stop cholera's spread: Professor proposes antibiotics for relief workers to prevent outbreaks

October 24 2012, by Alvin Powell



“The most likely conclusion is cholera was introduced in Haiti by a human,” said John Mekalanos, Lehman Professor of Microbiology and Molecular Genetics, in a talk called “River Monster: The Epidemiology, Ecology, and Pathobiology of Cholera.” Haiti had been cholera-free for 100 years before the devastating earthquake struck near Port-au-Prince in January 2010. Credit: Rose Lincoln/Harvard Staff Photographer

A Harvard medical specialist suggested Monday that relief workers and peacekeepers from cholera-endemic countries should be treated with antibiotics before they serve in other nations, to avoid repeating the Haitian epidemic that has killed thousands.

Haiti had been cholera-free for 100 years before the [devastating earthquake](#) struck near Port-au-Prince in January 2010. The disaster brought [relief workers](#) and United Nations' peacekeepers from around the world. The [cholera epidemic](#) began 10 months after the quake, with the bacterium possibly brought in by a Nepali peacekeeper. The outbreak has so far killed 7,000 and sickened 600,000 in Haiti, with 20,000 more cases in the nearby Dominican Republic.

"The most likely conclusion is cholera was introduced in Haiti by a human," said John Mekalanos, Lehman Professor of Microbiology and [Molecular Genetics](#) and head of Harvard Medical School's Department of Microbiology and Immunobiology.

Mekalanos delivered his comments at the Radcliffe Institute for Advanced Study's Fay House in a talk called "River Monster: The Epidemiology, Ecology, and Pathobiology of Cholera," part of the institute's water lecture series.

Cholera, which is caused by the [bacterium *Vibrio cholerae*](#), has been plaguing humans for millennia, with likely descriptions of it as far back as A.D. 40, Mekalanos said. The battle against cholera gave rise to epidemiology, which traces its roots to an 1854 London outbreak and physician John Snow's tracking the ailment to an infected water pump, whose handle city officials eventually removed.

Though a rarity in the developed world, cholera remains a scourge in developing nations. It infects an estimated 3 million to 5 million annually and kills 100,000, according to the U.S. Centers for Disease Control and

Prevention. Cholera, which causes vomiting and massive diarrhea, is easily treatable by modern medical science, but still deadly in places far removed from health care.

In addition, cholera appears to be getting more deadly, Mekalanos said. A new strain appeared in the 1960s and has slowly spread around the world. Another strain appeared in the 1990s and has begun spreading beyond its Asian birthplace. These two strains make up almost all the cholera in the world today, Mekalanos said.

Cholera is spread in water contaminated by disease-containing feces, which explains its absence in developed nations, with their advanced water and sewage treatment facilities. The disease is found in developing nations with poor sanitation. Once a person ingests a cholera bacterium, it multiplies in the intestines, releases cholera toxin, and then gets flushed out by the resulting diarrhea to infect new hosts.

Though there had been periodic cholera epidemics in the past, the New World had been cholera-free for about 100 years until 1991, when an epidemic began in Peru.

After the outbreak began in Haiti, Mekalanos began testing samples from Haiti and comparing them with samples in Peru and Bangladesh. He found that the Haitian strain was much more closely related to the Bangladesh strain, indicating it originated in Asia. A 2012 comparison with cholera strains from Nepal found an even closer match, supporting the contention that the bacteria arrived with Nepali peacekeepers near the outbreak site.

Mekalanos said little is known about cholera's seasonality and about what happens once it enters the environment. Research shows that some viruses prey on the bacteria while it is in the water, with the virus numbers negatively correlated to those of the cholera bacteria, rising as

cholera is falling. That raises the possibility of an environmental control of the disease while it is in the water supply.

"I'm humbled by how little we know [about [cholera](#)] in an ecological setting," Mekalanos said.

This story is published courtesy of the [Harvard Gazette](#), Harvard University's official newspaper. For additional university news, visit [Harvard.edu](#).

Provided by Harvard University

Citation: A plan to stop cholera's spread: Professor proposes antibiotics for relief workers to prevent outbreaks (2012, October 24) retrieved 3 May 2024 from <https://medicalxpress.com/news/2012-10-cholera-professor-antibiotics-relief-workers.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--