

# Report cites urgent need for revision of HIV prevention practices

October 17 2012

---



Trends in the sexual practices of gay men in Australia show that HIV prevention remains firmly embedded in gay communities. However, to curb the epidemic and start driving down numbers of new infections, "a re-invigoration of prevention approaches is now needed for Australia to continue its role as global leader in the HIV response," says Professor John de Wit, lead editor of the Annual Report of Trends in Behaviour 2012 (ARTB 2012).

The report draws on behavioural surveillance and research undertaken by the National Centre in HIV Social Research (NCHSR) at the University

of New South Wales (UNSW). The ARTB 2012 addresses changes in behaviour over time and identifies key emerging issues to deepen understanding of epidemiological trends and inform effective responses.

Findings from Gay Community Periodic Surveys (GCPS), reported in the 2012 ARTB, confirm that many [HIV prevention](#) practices remain well supported in the gay community. However, unprotected anal intercourse with casual partners, a main risk factor for [HIV infection](#), has gradually increased over the past decade. In 2011, one in three (34.6%) men who had sex with a casual partner reported unprotected anal intercourse in the past six months.

"It remains critical to continue to invest in effective programs and campaigns to reinforce condom use and inform men of the appropriate use of other sexual risk reduction practices, such as limiting condomless sex to partners of the same [HIV status](#)," said Professor de Wit.

The GCPS also shows that almost 90% of gay men have tested for HIV in their lifetime and 60% of HIV-negative gay men had tested in the past 12 months.

Professor de Wit said this shows the importance of making regular testing easier.

"Strengthening HIV testing facilities in community locations and the use of testing technologies that provide rapid results will be critical to further promote regular HIV testing, as is addressing the stigma that continues to be related to HIV, including in the gay community," he said.

The proportion of HIV-positive men in the GCPS nationally who were on combination antiretroviral treatment has slightly increased over the past decade from 66.2% in 2002 to 71.8% in 2011.

The proportion of HIV-positive participants reporting undetectable levels of virus in their blood has increased markedly from 52.6% in 2003 to 72.2% in 2011. Undetectable viral load that results from effective treatment indicates reduced infectiousness, but an online survey among [gay men](#) across Australia found that HIV-positive and HIV-negative men are sceptical about the preventive effects of HIV treatment.

The Annual Report of Trends in Behaviour 2012 will be released by UNSW's National Centre in HIV Social Research at the Australasian HIV/AIDS Conference 2012, Melbourne Convention and Exhibition Centre, Rooms 219 and 220. [hivaidsconference.com.au/](http://hivaidsconference.com.au/)

Provided by University of New South Wales

Citation: Report cites urgent need for revision of HIV prevention practices (2012, October 17) retrieved 25 April 2024 from <https://medicalxpress.com/news/2012-10-cites-urgent-hiv.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.