


Diverse forms of distress have distinct impact in diabetes

October 10 2012



In primary care patients with type 2 diabetes, depressive symptoms are predictive of future lifestyle-oriented self-management behaviors, while diabetes-related distress predicts glycemic control, possibly due to medication adherence, according to research published online Oct. 1 in *Diabetes Care*.

(HealthDay)—In primary care patients with type 2 diabetes, depressive symptoms (DS) are predictive of future lifestyle-oriented self-management behaviors, while diabetes-related distress (DRD) predicts glycemic control, possibly due to medication adherence, according to research published online Oct. 1 in *Diabetes Care*.

James E. Aikens, Ph.D., from the University of Michigan in Ann Arbor, conducted a longitudinal, six-month study involving 253 primary care patients with type 2 diabetes to compare DS and DRD as longitudinal predictors of medication adherence, self-care behavior, and glycemic control.

Aikens found that the results were similar with cross-sectional and longitudinal analyses. In longitudinal analyses, DS alone correlated significantly with future diet behavior, physical activity and [glucose testing](#). DRD alone was a significant predictor of future glycemic control and medication adherence.

"This study provides longitudinal support for the conceptual and empirical distinctions between DS and DRD in type 2 diabetes. DS may selectively suppress lifestyle-oriented self-management behaviors such as healthy eating, glucose testing, and exercise," Aikens concludes. "In contrast, diabetes-specific distress may impact subsequent [medication adherence](#) and glycemic control. Clinical assessment and intervention should encompass both factors."

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2012 [HealthDay](#). All rights reserved.

Citation: Diverse forms of distress have distinct impact in diabetes (2012, October 10) retrieved 10 April 2024 from
<https://medicalxpress.com/news/2012-10-diverse-distress-distinct-impact-diabetes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
