

# More GI bleeding seen in atrial fibrillation patients on rivaroxaban than warfarin

October 22 2012

---

Patients with atrial fibrillation (AF) experienced more major and non-major clinically relevant GI bleeding when taking rivaroxaban than patients taking warfarin.

As part of the ROCKET AF trial, researchers from multiple institutions, including Harvard Medical School; Mount Sinai Medical Center; Janssen Research & Development; and Bayer HealthCare Pharmaceuticals, randomized 14,264 patients with nonvalvular AF to receive either [rivaroxaban](#) or dose-adjusted [warfarin](#).

Post-hoc analysis was performed on all patients while receiving study medication until 2 days after the last dose.

Results showed that GI bleeding events (upper, lower, rectal) occurred more frequently in patients receiving rivaroxaban (n=394) than warfarin (n=290). In addition, major and nonmajor, clinically relevant bleeding occurred more frequently with rivaroxaban use than warfarin use.

However, there were fewer fatal GI bleeds with rivaroxaban and the absolute fatality rate was very low.

This study was presented during CHEST 2012, the annual meeting of the American College of Chest Physicians, held October 20 – 25, in Atlanta, Georgia.

Provided by American College of Chest Physicians

Citation: More GI bleeding seen in atrial fibrillation patients on rivaroxaban than warfarin (2012, October 22) retrieved 23 April 2024 from <https://medicalxpress.com/news/2012-10-gi-atrial-fibrillation-patients-rivaroxaban.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.