

Study shows group-based weight loss treatment effective whether led by health professionals or by peer counselors

October 9 2012



Study shows that group-based weight loss treatment is effective whether led by health professionals or by peer counselors. Credit: Baruch College

A new National Institutes of Health (NIH)-funded study conducted at Baruch College and published in *Obesity* found that overweight and obese adults who participated in three different weight loss treatments, all involving intensive, multi-component counseling delivered in groups, lost significant weight after 48 weeks whether the treatment was led by a health professional or by someone who had previous weight loss success.

This [randomized clinical trial](#) led by Angela Marinilli Pinto, PhD Assistant Professor of Psychology at Baruch College, is the first to compare professionally delivered group-based behavioral weight loss treatment— considered the "gold standard" for treating moderate [overweight and obesity](#)- to Weight Watchers, which is led by trained peer counselors who have achieved their own successful weight loss. The findings open up the possibility of offering effective, affordable weight loss options to more Americans.

In this study, 141 overweight and [obese men](#) and women were randomly assigned to:

- 48 weeks of behavioral weight loss treatment delivered by a health professional,
- 48 weeks of Weight Watchers, where group support is led by Weight Watchers members who have successfully achieved and maintained a healthy goal weight,
- Or a combined treatment consisting of 12 weeks of behavioral weight loss treatment delivered by a health professional followed by 36 weeks of Weight Watchers.

Weight Watchers was chosen because it is the largest commercial program in the United States and is behaviorally oriented and focuses on modifying diet and increasing [physical activity](#) to produce weight loss. The [investigators](#) hypothesized that the combined treatment would produce better weight losses than Weight Watchers alone. Assessments were conducted at baseline and weeks 12, 24, and 48, with weight change as the primary outcome.

All groups lost weight, but contrary to expectations, Weight Watchers participants lost more weight, on average, at the end of treatment compared to participants in the combination approach (mean weight loss

in the professionally led program was not significantly different from either group). In fact, 37% of Weight Watchers participants lost 10% or more of their starting weight compared to 15% of combined group participants and 11% of participants in the professionally led group.

Results of the study also showed that better meeting attendance was associated with greater weight losses during the first 24 weeks in all treatments. In the Weight Watchers group, this relationship was maintained even in the second half of treatment, suggesting that individuals who continue to stay involved in the Weight Watchers program do better.

"This study is important because, with almost 70% of American adults classified as overweight or obese, there is a need to provide practical treatment solutions that are effective, accessible, and affordable," Dr. Pinto explained. "Results of this study demonstrate that the Weight Watchers program can produce clinically meaningful [weight](#) losses." The cost of joining [Weight Watchers](#) is approximately \$10 per week, which includes member registration, weekly meetings, and access to online resources.

More information: A.M. Pinto et al: Combining behavioral weight loss treatment and a commercial program: a randomized clinical trial. *Obesity* 2012; [DOI: 10.1002/oby.20044](https://doi.org/10.1002/oby.20044)

Provided by Baruch College

Citation: Study shows group-based weight loss treatment effective whether led by health professionals or by peer counselors (2012, October 9) retrieved 5 May 2024 from <https://medicalxpress.com/news/2012-10-group-based-weight-loss-treatment-effective.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.