

Sitting for protracted periods increases risk of diabetes, heart disease and death

October 15 2012

A new study led by the University of Leicester, in association with colleagues at Loughborough University, has discovered that sitting for long periods increases your risk of diabetes, heart disease and death.

The study, which combined the results of 18 studies and included a total of 794,577 participants, was led by Dr. Emma Wilmot, a research fellow in the [Diabetes](#) Research Group at the University of Leicester. It was done in collaboration with colleagues from the newly established National Institute for [Health Research](#) (NIHR) Leicester-Loughborough Diet, Lifestyle and Physical Activity Biomedical Research Unit and was published in *Diabetologia*, the journal of the European Association of the Study of Diabetes.

According to the study, those who sit for long periods have a two fold increase in their risk of diabetes, [heart disease](#) and death. Importantly, associations were independent of the amount of moderate-to-[vigorous physical activity](#) undertaken, suggesting that even if an individual meets typical [physical activity guidelines](#), their health may still be compromised if they sit for long periods of time throughout the day.

Dr Wilmot, a Clinical Research Fellow in Diabetes and Endocrinology based at the Leicester Diabetes Centre, Leicester General Hospital, said: "The average adult spends 50-70% of their time sitting so the findings of this study have far reaching implications. By simply limiting the time that we spend sitting, we may be able to reduce our risk of diabetes, heart disease and death".

"Our study also showed that the most consistent associations were between sitting and diabetes. This is an important message because people with risk factors for diabetes, such as the obese, those of South Asian [ethnic origin](#), or those with a family history of diabetes, may be able to help reduce their future risk of diabetes by limiting the time spent sitting. "

Professor Stuart Biddle, of Loughborough University, and a co-investigator on the study, said: "There are many ways we can reduce our sitting time, such as breaking up long periods at the computer at work by placing our laptop on a filing cabinet. We can have standing meetings, we can walk during the lunch break, and we can look to reduce TV viewing in the evenings by seeking out less sedentary behaviours."

Professor Melanie Davies, Professor of Diabetes Medicine at the University of Leicester and honorary consultant at University Hospitals of Leicester is a co-investigator and Director of the NIHR Leicester-Loughborough Diet, Lifestyle and Physical Activity [Biomedical Research](#) Unit. She said:

"This paper has a very important message for the public but also for health care professionals - namely that being sedentary is common and dangerous for our long term health, particularly for diabetes and cardiovascular disease, and that this link appears to be over and above other lifestyle factors such as our diet and [physical activity](#)."

Provided by University of Leicester

Citation: Sitting for protracted periods increases risk of diabetes, heart disease and death (2012, October 15) retrieved 5 May 2024 from <https://medicalxpress.com/news/2012-10-protracted-periods-diabetes-heart-disease.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.