

Bariatric surgical procedures have similar therapeutic benefits in obese adults

November 26 2012

Obesity is associated with insulin resistance and type 2 diabetes, both of which can be significantly improved by weight loss. Gastric bypass and adjustable gastric banding are two bariatric surgery techniques that are frequently used to effect weight loss in obese patients, but it is unclear if the two procedures produce different outcomes.

In this issue of the [Journal of Clinical Investigation](#), researchers led by Samuel Klein at the University of Washington School of Medicine in St. Louis compared the effects of 20% weight loss induced by either gastric bypass or adjustable gastric banding on metabolic response.

They found that patients had different metabolic responses after eating, but both procedures equally improved insulin sensitivity and [glucose tolerance](#).

The researchers concluded that weight loss itself is primarily responsible for the therapeutic effects of gastric bypass and adjustable gastric banding in non-diabetic obese adults.

More information: Gastric bypass and banding equally improve insulin sensitivity and β -cell function, *Journal of Clinical Investigation*, 2012.

Provided by Journal of Clinical Investigation

Citation: Bariatric surgical procedures have similar therapeutic benefits in obese adults (2012, November 26) retrieved 23 April 2024 from <https://medicalxpress.com/news/2012-11-bariatric-surgical-procedures-similar-therapeutic.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.