

# Novel breast screening technology increases diagnostic accuracy

November 20 2012

---

The addition of three-dimensional breast imaging—a technology called tomosynthesis—to standard digital mammography significantly increases radiologists' diagnostic accuracy while reducing false positive recall rates, according to the results of a multi-center study published in *Radiology*.

"This is the first major advance in breast imaging and [breast cancer screening](#) since the development of [breast MRI](#)," said lead researcher Elizabeth A. Rafferty, M.D., director of Breast Imaging at the Avon Comprehensive Breast Center at Massachusetts General Hospital in Boston. "The beauty of tomosynthesis is that it addresses two major concerns with screening mammography: missed cancers and false positive rates."

According to the [National Cancer Institute](#), there is convincing evidence that [screening mammography](#) reduces [breast cancer mortality](#) in women between the ages of 40 and 74. However, as many as 30 percent of breast cancers are not detected by mammography and an additional eight to 10 percent of women who undergo a [screening mammogram](#) are recalled for further testing when no cancer is present (called a false positive result).

Unlike a screening digital mammogram, which involves two X-ray images of each breast, breast tomosynthesis captures multiple, low-dose images from different angles around the breast. The images are then used to produce a three-dimensional reconstruction of the breast. Both

[digital mammography](#) and breast tomosynthesis, which was approved by the United States Food and Drug Administration (FDA) in February of 2011, can be performed on the same mammography equipment in rapid succession.

Dr. Rafferty's study involved 1,192 women recruited from five sites, of whom 997, including 780 screening cases and 217 women who needed pre-biopsy [breast imaging](#), had complete data sets. Each of the women underwent a standard digital mammogram followed by breast tomosynthesis. The total radiation dose for the combined procedure was less than 3 milligray, which is the FDA limit for a single mammogram.

Drawing from the eligible cases, the researchers then conducted two reader studies involving 312 and 310 cases, respectively. Twelve radiologists participated in the first reader study; 15 radiologists in the second. A total of 48 cancers were included in the first reader study; 51 cancers in the second.

Compared to digital mammography alone, the use of both standard mammogram and tomosynthesis resulted in increased diagnostic accuracy for all 27 radiologists. Additionally, the diagnostic sensitivity of the combined exam—or the rate at which cancer present in the breast was correctly identified—increased by 10.7 percent for radiologists in Reader Study 1 and 16 percent for radiologists in Reader Study 2.

"Almost all of the gains in diagnostic sensitivity with the combined modality were attributable to the improved detection and characterization of invasive cancers, which are the cancers we are most concerned about because of their potential to metastasize," Dr. Rafferty said.

With the addition of breast tomosynthesis to standard digital mammography, false positive recall rates also significantly decreased for

all 27 of the radiologists. Absolute recall rate reductions of 38.6 percent and 17.1 percent were seen in Reader Studies 1 and 2, respectively.

"In the clinical setting, we would expect that type of reduction in recall rate to translate into a substantial number of unnecessary diagnostic tests being avoided," Dr. Rafferty said.

**More information:** "Assessing Radiologist Performance Using Combined Digital Mammography and Breast Tomosynthesis Compared to Digital Mammography Alone: Results of a Multicenter, Multireader Trial." *Radiology*, 2012.

Provided by Radiological Society of North America

Citation: Novel breast screening technology increases diagnostic accuracy (2012, November 20) retrieved 25 April 2024 from

<https://medicalxpress.com/news/2012-11-breast-screening-technology-diagnostic-accuracy.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.