

Chronic fatigue syndrome—a system under stress

November 15 2012

Australian researchers have discovered for the first time that reduced heart rate variability – or changes in heart beat timing – best predicts cognitive disturbances, such as concentration difficulties commonly reported by people with chronic fatigue syndrome (CFS). This adds to the growing body of evidence linking autonomic nervous system imbalance to symptoms of this poorly understood disorder.

The findings are reported in the journal [PLOS ONE](#).

[Chronic fatigue syndrome](#) is characterised by medically unexplained, disabling fatigue and [neuropsychiatric symptoms](#) of at least six months' duration. The disturbance underlying the symptoms in CFS is still poorly understood.

"We have studied autonomic function in CFS for some time and our findings clearly indicate a loss of integrity in stress-responsive neural and physiological systems in CFS. Patients with this condition are hyper-responsive to challenges arising both from within the body and from the environment," says lead researcher, Associate Professor Ute Vollmer-Conna of the University of New South Wales in Sydney, Australia.

"Even when they sleep, their stress-responsive neural systems are on high alert, signalling that it is not safe to relax. I think this condition may be understood by analogy to post-traumatic stress disorder, just that in CFS the original trauma is most likely a physiological, internal one, such as a severe infection."

In a study of 30 patients with CFS and 40 healthy individuals, UNSW researchers recorded the heart beats of participants (via ECG) and analysed cardiac responses to cognitive challenges, and associations with [mental performance](#) outcomes.

The patients with CFS performed with similar accuracy, but they took significantly longer to complete the tests than people without the condition. They also had greater heart rate reactivity; low and unresponsive [heart rate variability](#); and prolonged heart rate-recovery after the cognitive challenge.

[Resting heart rate](#) variability (an index of vagus nerve activity) was identified as the only significant predictor of cognitive outcomes, while current levels of fatigue and other symptoms did not relate to cognitive performance.

"This is the first demonstration of an association between reduced cardiac vagal tone and cognitive impairment in CFS. Our findings confirm previous reports of a significant loss of vagal modulation, which becomes particularly apparent when dealing with challenging tasks. The current results are consistent with the notion that CFS represents a 'system under stress'," Associate Professor Vollmer-Conna says.

The findings could lead to new ways to improve cognitive difficulties in people with CFS, including biofeedback assisted retraining of autonomic functioning, the researchers say.

More information: [dx.plos.org/10.1371/journal.pone.0049518](https://doi.org/10.1371/journal.pone.0049518)

Provided by University of New South Wales

Citation: Chronic fatigue syndrome—a system under stress (2012, November 15) retrieved 19 April 2024 from

<https://medicalxpress.com/news/2012-11-chronic-fatigue-syndrome-a-stress.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.