

# Older adults who are frail much more likely to be food insufficient, according to national study

November 5 2012

---

A national study of older Americans shows those who have limited mobility and low physical activity – scientifically categorized as "frail" – are five times more likely to report that they often don't have enough to eat, defined as "food insufficiency," than older adults who were not frail.

The nationally representative study of more than 4,700 adults older than age 60 in the United States uses data from the Third National Health and [Nutrition Examination Survey](#). The results are online today in the [British Journal of Nutrition](#).

Lead author Ellen Smit, an epidemiologist at Oregon State University, said food insufficiency occurs when people report that they sometimes or often do not have enough food to eat. Food-insufficient older adults have been shown to have poor [dietary intake](#), [nutritional status](#) and health status.

"Although little is known about food insufficiency as it relates to frailty, conceivably we thought if food insufficiency is associated with poorer nutritional status, it may also be associated with physical functioning and frailty," she said.

Frailty is a state of decreased physical functioning and a significant complication of aging that increases the risk for incident falls, fractures, disability, [health care expenditures](#), and [premature mortality](#). People in

this study are diagnosed as frail when they meet two of the following criteria: slow walking, muscular weakness, exhaustion and low physical activity.

Smit said as the population ages, with more than 20 percent of Americans expected to be older than 65 by 2030, the need for identifying clinical and population-based strategies to decrease the prevalence and consequences of frailty are needed. In her study, almost 50 percent of people were either frail, or "pre-frail," meaning that they were at risk for decreased physical functioning.

Frail people were older, less educated, at lower income levels, more likely to be female, more likely to be smokers, and less likely to be white than adults who were not frail. Frail people were also more likely to be either underweight or obese, while at the same time eating fewer calories than people who were not frail.

"We need to target interventions on promoting availability and access to nutritious foods among frail [older adults](#)," Smit said. "It is also important to improve nutritional status while not necessarily increasing body weight."

Frail adults may have difficulty leaving the house, for instance, and accessing fresh fruits and vegetables. Smit said communities could work on identifying programs or nonprofit organizations that can deliver nutritious meals or fresh produce to older frail adults.

Provided by Oregon State University

Citation: Older adults who are frail much more likely to be food insufficient, according to national study (2012, November 5) retrieved 26 April 2024 from <https://medicalxpress.com/news/2012-11-older-adults-frail-food-insufficient.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.