


Statins plus exercise best at lowering cholesterol, study finds

November 28 2012, by Steven Reinberg, Healthday Reporter



Moderate exercise reduced death risk even without drugs, but not as much.

(HealthDay)—People who exercise along with taking statins to lower their high cholesterol levels can dramatically reduce their risk of dying, a large new study suggests.

"The reduction in death is independent; whatever statins do is independent of what exercise does," said lead researcher Peter Kokkinos, a professor in the cardiology department at the Veterans Affairs Medical Center in Washington, D.C.

"When you combine the two, you get even better results," he said. "If you are taking statins, your mortality is about 35 percent lower versus not taking statins, but if you exercise, your mortality level decreases as your fitness level increases to the point where you can reach a 70 percent

reduction in mortality."

Kokkinos is talking about regular [moderate exercise](#)—not vigorous workouts. "Thirty minutes a day of brisk walking—not a whole lot," he said.

[Statin drugs](#) include Lipitor ([atorvastatin](#)), Lescol (fluvastatin), Pravachol ([pravastatin](#)), Crestor ([rosuvastatin](#)) and Zocor ([simvastatin](#)).

Some people can't take statins because of side effects, Kokkinos said. "For these people, exercise alone reduces your risk just as much, if not more, than statins," he said. However, he stressed, "we do not recommend that people do not take their statins."

Exercise works by stressing the body making it stronger, Kokkinos said. It's an [evolutionary adaptation](#) to protect the body from being overcome by changing stressors, he said.

"Get off the couch—walk," Kokkinos said. "About 150 minutes a week of brisk walking is all you need."

The report was published online in the Nov. 28 edition of *The Lancet*.

For the study, Kokkinos' team analyzed the medical records of more than 10,000 veterans with [high cholesterol levels](#) treated in Veterans Affairs hospitals in Washington D.C., and Palo Alto, Calif. Of these, 9,700 were men and 343 were women.

The researchers judged the fitness levels of the participants by looking at the results of standard [treadmill exercise](#) tolerance tests, which were given between 1986 and 2011.

Fewer deaths occurred among participants who were taking statins and

were physically fit. Over 10 years of follow-up, those who were the most physically fit had the lowest risk of dying, the researchers noted.

These fittest people actually reduced their risk of dying by about 60 percent regardless of whether they were taking statins, according to the study.

These difference in death risk could not be explained by age, weight, ethnicity, gender, history of cardiovascular disease, risk factors for cardiovascular disease or medications, the researchers said.

According to the U.S. Centers for Disease Control and Prevention, about 71 million Americans have high cholesterol, which is an important risk factor for heart disease, the authors pointed out.

One expert familiar with the new findings cautioned that exercise is not a replacement for statins in those with high cholesterol. The best results are among those taking statins and who are the fittest, he emphasized.

"Cardiovascular disease remains the leading cause of death and disability in men and women in the United States," said Dr. Gregg Fonarow, a spokesman for the American Heart Association and a professor of cardiology at the University of California, Los Angeles.

"Statin therapy has been proven in multiple clinical trials to substantially reduce cardiovascular events and all-cause mortality in men and women with or at risk for cardiovascular disease," he said.

Some people have assumed that if they are engaging in regular exercise and are physically fit that they may not benefit from statin treatment and some physicians consider statin therapy only for patients who have failed attempts at lifestyle modification, Fonarow said.

This new study shows that at all levels of physical fitness, statin therapy was independently associated with lower risk of dying. The very best 10-year outcomes were among men and women taking [statins](#) with the highest [fitness levels](#), he noted.

"These findings further reinforce the remarkable real-world clinical effectiveness and safety of statin therapy to prevent and treat [cardiovascular disease](#)," Fonarow said.

More information: To learn more about cholesterol, visit the [American Heart Association](#).

Copyright © 2012 [HealthDay](#). All rights reserved.

Citation: Statins plus exercise best at lowering cholesterol, study finds (2012, November 28) retrieved 1 May 2024 from <https://medicalxpress.com/news/2012-11-statins-lowering-cholesterol.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--