


# New tools reveal 'new beginning' in split-brain research

November 2 2012



This diagram shows changes in the interhemispheric network over time. The top row (blue) shows the interhemispheric network when a stimulus was shown to the right brain (left visual field). The bottom row (orange) shows the interhemispheric network when a stimulus was shown to the left brain (right visual field). More network connections are seen when the stimulus was shown to the right brain and transferred to the left (top row). Credit: UCSB

(Medical Xpress)—UC Santa Barbara has reported an important discovery in the interdisciplinary study of split-brain research. The findings uncover dynamic changes in brain coordination patterns

between left and right hemispheres.

Split-brain research has been conducted for decades, and scientists have long ago shown that language processing is largely located in the left side of the brain. When words appear only in the left visual field—an area processed by the right side of the brain—the right brain must transfer that information to the left brain, in order to interpret it. The new study at UC Santa Barbara shows that healthy test subjects respond less accurately when information is shown only to the right brain.

While hemispheric specialization is considered accurate, the new study sheds light on the highly complex interplay—with neurons firing back and forth between distinct areas in each half of the brain. The findings rely on extremely sensitive neuroscience equipment and analysis techniques from network science, a fast-growing field that draws on insights from sociology, mathematics, and physics to understand complex systems composed of many interacting parts. These tools can be applied to systems as diverse as earthquakes and brains.

Fifty years ago, UC Santa Barbara neuroscientist Michael S. Gazzaniga moved the field forward when he was a graduate student at the California Institute of Technology and first author of a groundbreaking report on split-brain patients. The study, which became world-renowned, was published in the [Proceedings of the National Academy of Sciences](#) (PNAS) in August 1962. This week, in the very same journal, Gazzaniga and his team announced major new findings in split-brain research. The report is an example of the interdisciplinary science for which UC Santa Barbara is well known.

"The occasion of this paper is on the 50th anniversary of the first report on human split-[brain research](#) reported in PNAS," said Gazzaniga. "That study showed how surgically dividing the two hemispheres of the human brain—in an attempt to control epilepsy—allowed for studying how each

isolated half-brain was specialized for cognitive function.

"In the present study, new techniques—not present 50 years ago—begin to allow for an understanding of how the normal, undivided brain integrates the special functions of each half brain. It is a new beginning and very exciting," said Gazzaniga, professor of psychology in UC Santa Barbara's Department of Psychological and Brain Sciences, and director of the university's SAGE Center for the Study of Mind.

Karl W. Doron, first author and a postdoctoral scholar with Gazzaniga, said: "This study shows that the strength of communication across the brain changes dynamically, when information is exchanged between the right and left hemispheres via the corpus callosum, which connects both sides of the brain. The findings shed light on the neural underpinnings of inter-hemispheric communication first reported 50 years ago in the classic split-brain studies performed by Gazzaniga and colleagues."

Doron explained that years of research—in both clinical patient populations and healthy individuals—have shown that the brain is made up of specialized modules that preferentially process certain types of information.

When linguistic information, such as a word or pronounceable non-word, is shown directly to the left brain, people are better at deciding whether or not it is a real word, said Doron.

"In contrast, when the information is shown directly to the right brain, people perform worse," he said. "It has long been assumed that language information shown to the right brain must be transferred through the corpus collosum. In other words, right brain information must be directly transferred to the language-specialized left brain, in order to be processed correctly."

In the new study, researchers used magnetoencephalography, or MEG, a non-invasive tool that measures the continuous brain activity of nine healthy young adults. The participants were briefly flashed a stimulus that was either a real word or a "word-like" non-word. The stimulus was randomly shown either to the left or right brain, using a split-visual field experimental technique.

MEG allows researchers to peer into brain activity at the rate of 512 images per second, while fMRI only allows for images to be recorded at the rate of one image every two seconds. The researchers processed this temporal data using novel network analysis techniques drawn from the mathematical and physical sciences that enable them to examine the dynamic nature of brain communication. This novel network analysis allows the data to be presented in movie form, rather than as snapshots. The researchers found that the strength of connections between the left and right hemispheres depended on whether the stimulus was shown to the left or right brain hemisphere.

"A larger number of statistically significant connections were present at several time points after the stimulus when the word or pronounceable non-word was shown to the right brain and needed to be transferred to the language-specialized left brain," said Danielle S. Bassett, UC Santa Barbara Sage Junior Research Fellow in the Department of Physics and the Department of Psychological and Brain Sciences.

"The results further suggest that brain function measured with MEG can be broken into functional communities that rearrange their organization over time — based on whether the left or right brain saw the stimulus, and thus whether inter-hemispheric information transfer must take place," said Bassett.

While working on his Ph.D. with Gazzaniga, which he completed in 2011, Doron directed the MEG experiments at the Otto-von-Guericke

University in Magdeburg, Germany, over a two-month period, with support from Gazzaniga's German Humboldt Fellowship Award. The MEG machine is composed of a vertical cylinder that is positioned over the head and held in place with a type of helmet.

Doron's research combines data from cognitive experiments with a variety of neuroimaging modalities to understanding the neuroscientific underpinnings of the mind, with specific focus on neural oscillatory processes and attention. Doron's current work merges graph theoretical, information theoretical, and complex systems approaches in an effort to describe the large-scale functional connectivity of [brain](#) networks.

Provided by University of California - Santa Barbara

Citation: New tools reveal 'new beginning' in split-brain research (2012, November 2) retrieved 6 May 2024 from <https://medicalxpress.com/news/2012-11-tools-reveal-split-brain.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--