

ASH: More intense treatment beneficial in polycythemia vera

December 10 2012



For patients with polycythemia vera, more intense treatment to achieve a hematocrit target of less than 45 percent correlates with a significantly lower rate of cardiovascular death and major thrombosis compared with a target of 45 to 50 percent, according to a study published online Dec. 8 in the *New England Journal of Medicine* to coincide with presentation at the annual meeting of the American Society of Hematology, held from Dec. 8 to 11 in Atlanta.

(HealthDay)—For patients with polycythemia vera, more intense treatment to achieve a hematocrit target of less than 45 percent correlates with a significantly lower rate of cardiovascular death and major thrombosis compared with a target of 45 to 50 percent, according to a study published online Dec. 8 in the *New England Journal of Medicine* to coincide with presentation at the annual meeting of the American Society of Hematology, held from Dec. 8 to 11 in Atlanta.

Roberto Marchioli, M.D., from the Consorzio Mario Negri Sud in Santa Maria Imbaro, Italy, and colleagues conducted a <u>randomized controlled</u>



trial involving 365 adults with JAK2-positive polycythemia vera to compare more intensive treatment (target hematocrit less than 45 percent; low-hematocrit group) or less intensive treatment (target hematocrit 45 to 50 percent; high-hematocrit group) with phlebotomy, <u>hydroxyurea</u>, or both.

After a median follow-up of 31 months, the researchers found that the primary end point (time until death from cardiovascular causes or major thrombotic events) was reached by 2.7 percent of patients in the low-hematocrit group and 9.8 percent in the high-hematocrit group (hazard ratio in high-hematocrit group, 3.91). The primary end point plus superficial-vein thrombosis was reached by 4.4 and 10.9 percent, respectively (hazard ratio, 2.69). The rates of adverse events were similar between the groups.

"In conclusion, among patients with polycythemia vera, maintaining a hematocrit target of less than 45 percent, as compared with a target of 45 to 50 percent, was associated with a significantly lower rate of thrombotic complications without an increase in serious treatment complications," the authors write.

Several authors disclosed financial ties to the pharmaceutical industry.

More information: <u>Abstract</u> <u>Full Text</u> <u>Editorial</u> <u>More Information</u>

Copyright © 2012 <u>HealthDay</u>. All rights reserved.

Citation: ASH: More intense treatment beneficial in polycythemia vera (2012, December 10) retrieved 2 May 2024 from



https://medicalxpress.com/news/2012-12-ash-intense-treatment-beneficial-polycythemia.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.