

Host cholesterol secretion likely to influence gut microbiota

December 18 2012

For more than half a century, researchers have known that the bacteria that colonize the gastrointestinal tract of mammals influence their host's cholesterol metabolism. Now, Jens Walter and colleagues of the University of Nebraska show that changes in cholesterol metabolism induced by diet can alter the gut flora. The research was published online ahead of print in the journal *Applied and Environmental Microbiology*.

In the study, the researchers added plant sterol esters to the diets of hamsters. The overall effect of this was to inhibit several bacterial taxa, from the families Coriobacteriacea and Erysipelotrichaceae, says Walter. But the immediate effect of the plant sterols was to physically block cholesterol absorption by the intestine. That decreased <u>cholesterol levels</u> in the liver and the plasma, prompting the hamster's body to respond by synthesizing more cholesterol. That, in turn, boosted cholesterol excretion into the gut, and that extra cholesterol was the direct inhibitor of those bacterial families.

"The abundance of these bacterial taxa and the levels of cholesterol in the fecal samples followed a mathematical model of bacterial inhibition," says Walter.

Practically speaking, the microbial inhabitants of the gut are part of the <u>metabolic system</u>. Researchers have shown that certain health problems are related to changes in the <u>gut flora</u>, such as can be induced by <u>overuse</u> <u>of antibiotics</u>. Since changes in diet can influence composition of the gut flora, health problems such as obesity might be targeted by dietary



interventions designed to suppress bacteria that contribute to weight gain. "However, for these to be successful, we need to know which bacterial patterns not only are associated with disease, but actually contribute to it," says Walter, noting that his research showed that some alterations associated with metabolic disease might be the consequence, rather than the cause of the disorder.

Walter says that the work was a real student project. Among the coauthors, three were graduate students, and two were undergraduates. "As a supervisor, it is extremely nice to see young scientists work as a team, and staying dedicated through the five years that this project took to complete," he says.

More information: A copy of the manuscript can be found online at <u>bit.ly/asmtip1212b</u>. Formal publication is scheduled for the January 2013 issue of *Applied and Environmental Microbiology*.

I. Martinez, D.J. Perdicaro, A.W. Brown, S. Hammons, T.J. Carden, T.P. Carr, K.M. Eskridge, and J. Walter, 2012. Diet-induced alterations of host cholesterol metabolism are likely to affect gut microbiota composition in hamsters. *Appl. Environ. Microbiol.* 2012 Nov 2. [Epub ahead of print]

Provided by American Society for Microbiology

Citation: Host cholesterol secretion likely to influence gut microbiota (2012, December 18) retrieved 28 April 2024 from <u>https://medicalxpress.com/news/2012-12-host-cholesterol-secretion-gut-microbiota.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.