

# Regular marijuana use by teens continues to be a concern

December 19 2012

---

Continued high use of marijuana by the nation's eighth, 10th and 12th graders combined with a drop in perceptions of its potential harms was revealed in this year's Monitoring the Future survey, an annual survey of eighth, 10th, and 12th-graders conducted by researchers at the University of Michigan. The survey was carried out in classrooms around the country earlier this year, under a grant from the National Institute on Drug Abuse (NIDA), part of the National Institutes of Health.

The 2012 survey shows that 6.5 percent of high school seniors smoke marijuana daily, up from 5.1 percent five years ago. Nearly 23 percent say they smoked it in the month prior to the survey, and just over 36 percent say they smoked within the [previous year](#). For 10th graders, 3.5 percent said they use marijuana daily, with 17 percent reporting past month use and 28 percent reporting use in the past year. The use escalates after eighth grade, when only 1.1 percent reported daily use, and 6.5 percent reported past month use. More than 11 percent of eighth graders said they used marijuana in the past year.

The Monitoring the Future survey also showed that teens' perception of marijuana's harmfulness is down, which can signal future increases in use. Only 41.7 percent of eighth graders see occasional use of marijuana as harmful; 66.9 percent see regular use as harmful. Both rates are at the lowest since the survey began tracking [risk perception](#) for this age group in 1991. As teens get older, their [perception](#) of risk diminishes. Only 20.6 percent of 12th graders see occasional use as harmful (the lowest

since 1983), and 44.1 percent see regular use as harmful, the lowest since 1979.

A 38-year NIH-funded study, published this year in the [Proceedings of the National Academy of Sciences](#), showed that people who used cannabis heavily in their teens and continued through adulthood showed a significant drop in [IQ](#) between the ages of 13 and 38—an average of eight points for those who met criteria for cannabis dependence. Those who used marijuana heavily before age 18 (when the brain is still developing) showed impaired mental abilities even after they quit taking the drug. These findings are consistent with other studies showing a link between prolonged marijuana use and cognitive or neural impairment.

"We are increasingly concerned that regular or daily use of marijuana is robbing many young people of their potential to achieve and excel in school or other aspects of life," said NIDA Director Nora D. Volkow, M.D. "THC, a key ingredient in marijuana, alters the ability of the hippocampus, a brain area related to learning and memory, to communicate effectively with other brain regions. In addition, we know from recent research that marijuana use that begins during adolescence can lower IQ and impair other measures of mental function into [adulthood](#)."

Research clearly demonstrates that marijuana has the potential to cause problems in daily life or make a person's existing problems worse. In one study, heavy marijuana abusers reported that the drug impaired several important measures of well-being and life achievement, including physical and mental health, cognitive abilities, social life, and career status.

"We should also point out that marijuana use that begins in adolescence increases the risk they will become addicted to the drug," said Volkow. "The risk of addiction goes from about 1 in 11 overall to about 1 in 6 for

those who start using in their teens, and even higher among daily smokers."

Use of other illicit drugs among teens continued a steady modest decline. For example, past year illicit drug use (excluding marijuana) was at its lowest level for all three grades at 5.5 percent for eighth graders, 10.8 percent for 10th graders, and 17 percent for 12th graders. Among the most promising trends, the past year use of Ecstasy among seniors was at 3.8 percent, down from 5.3 percent last year.

"Each new generation of young people deserves the chance to achieve its full potential, unencumbered by the obstacles placed in the way by drug use," said Gil Kerlikowske, director of National Drug Control Policy. "These long-term declines in youth drug use in America are proof that positive social change is possible. But now more than ever we need parents and other adult influencers to step up and have direct conversations with young people about the importance of making healthy decisions. Their futures depend on it."

The survey also looks at abuse of drugs that are easily available to teens because they are generally legal, sometimes for adults only (tobacco and alcohol), for other purposes (over-the-counter or prescribed medications; inhalants), or because they are new drugs that have not yet been banned. Most of the top drugs or drug classes abused by 12th graders are legally accessible, and therefore easily available to teens.

For the first time, the survey this year measured teen use of the much publicized emerging family of drugs known as "bath salts," containing an amphetamine-like stimulant that is often sold in drug paraphernalia stores. The data showed a relative low use among 12th graders at 1.3 percent. In addition, the survey measured use of the hallucinogenic herb Salvia, finding that past year use dropped among 10th and 12th graders, down to 4.4 percent for 12th graders from last year's 5.9 percent.

Abuse of synthetic marijuana (also known as K-2 or Spice) stayed stable in 2012 at just over 11 percent for past year use among 12th graders. While many of the ingredients in Spice have been banned by the U.S. Drug Enforcement Administration, manufacturers attempt to evade these legal restrictions by substituting different chemicals in their mixtures. Another drug type – inhalants – continues a downward trend. As one of the drugs most commonly used by younger students, the survey showed a past year use rate of 6.2 percent among eighth graders, a significant drop in the last five years when the 2007 survey showed a rate of 8.3 percent.

The data shows a mixed report regarding prescription drug abuse. Twelfth graders reported non-medical use of the opioid painkiller Vicodin at a past year rate of 7.5 percent. Since the survey started measuring its use in 2002, rates hovered near 10 percent until 2010, when the survey started reporting a modest decline. However, past year abuse of the stimulant Adderall, often prescribed to treat ADHD, has increased over the past few years to 7.6 percent among high school seniors, up from 5.4 percent in 2009. Accompanying this increased use is a decrease in the perceived harm associated with using the drug, which dropped nearly 6 percent in the past year—only 35 percent of 12th graders believe that using Adderall occasionally is risky. The survey continues to show that most teens who abused prescription medications were getting them from family members and friends.

The survey also measured abuse of over-the-counter cough and cold medicines containing dextromethorphan—5.6 percent of [high school seniors](#) abused them in the past year, a rate that has held relatively steady over the past five years.

The 2012 results also showed a continued steady decline in alcohol use, with reported use at its lowest since the survey began measuring rates. More than 29 percent of eighth graders said they have used alcohol in

their lifetime, down from 33.1 percent last year, and significantly lower than that peak rate of 55.8 percent in 1994. For 10th graders, 54 percent of teens reported lifetime use of alcohol, down from its peak of 72 percent in 1997. Binge drinking rates (five or more drinks in a row in the previous two weeks) have been slowly declining for eighth graders, at 5.1 percent, down from 6.4 percent in 2011, and 13.3 percent at their peak in 1996.

Cigarette smoking continues at its lowest levels among eighth, 10th and 12th graders, with dramatic long-term improvement. Significant declines were seen in lifetime use among eighth graders, down to 15.5 percent from last year's 18.4 percent, compared to nearly 50 percent at its peak in 1996. Significant declines were also seen in 10th grade lifetime use of cigarettes, down to 27.7 percent from 30.4 percent in 2011. Peak rates for 10th graders were seen in 1996 at 61.2 percent. For some indicators, including past month use in all three grades, cigarette smoking remains lower than marijuana use, a phenomenon that began a few years ago.

The [survey](#) also measures several other kinds of tobacco delivery products. For example, past year use of small cigars was reported at nearly 20 percent for 12th graders, with an 18.3 percent rate for hookah water pipes.

"We are very encouraged by the marked declines in tobacco use among youth. However, the documented use of non-cigarette tobacco products continues to be a concern," said Howard K. Koh, M.D., M.P.H., assistant secretary for health for the U.S. Department of Health and Human Services. "Preventing addiction includes helping kids be tobacco free so they can enjoy a fighting chance for health."

Provided by National Institutes of Health

Citation: Regular marijuana use by teens continues to be a concern (2012, December 19)  
retrieved 26 April 2024 from  
<https://medicalxpress.com/news/2012-12-regular-marijuana-teens.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.