

Detrimental effect of obesity on lesions associated with Alzheimer's disease

January 7 2013

Researchers from Inserm and the Université Lille/Université Lille Nord de France have recently used a neurodegeneration model of Alzheimer's disease to provide experimental evidence of the relationship between obesity and disorders linked to the tau protein. This research was conducted on mice and is published in the *Diabetes Review*. It corroborates the theory that metabolic anomalies contribute massively to the development of dementia.

In France, more than 860,000 people suffer from Alzheimer's disease and related disorders, making them the largest cause of age-related loss of intellectual function. Cognitive impairments observed in Alzheimer's disease result from the accumulation of abnormal tau proteins in nerve cells undergoing degeneration. We know that obesity, a major risk factor in the development of insulin resistance and type 2 diabetes, increases the risk of dementia during the aging process. However, the effects of obesity on 'Taupathies' (i.e. tau protein-related disorders), including Alzheimer's disease, were not clearly understood. In particular, researchers assumed that insulin resistance played a major role in terms of the effects of obesity.

The "Alzheimer & Tauopathies" team from mixed research unit 837 (Inserm/Université Lille 2/Université Lille Nord de France) directed by Dr. Luc Buée, in collaboration with mixed research unit 1011 "Nuclear receptors, cardiovascular diseases and diabetes", have just demonstrated, in mice, that obese subjects develop aggravated disorders. To achieve this result, young transgenic mice, who develop tau-related



neurodegeneration progressively with age, were put on a high-fat diet for five months, leading to progressive obesity.

"At the end of this diet, the obese mice had developed an aggravated disorder both from the point of view of memory and modifications to the Tau protein" explains David Blum, in charge of research at Inserm.

This study uses a neurodenegeneration model of Alzheimer's disease to provide <u>experimental evidence</u> of the relationship between obesity and disorders linked to the tau protein. Furthermore, it indicates that insulin resistance is not the aggravating factor, as was suggested in previous studies.

"Our research supports the theory that environmental factors contribute massively to the development of this neurodegenerative disorder" underlines the researcher. "Our work is now focussing on identifying the factors responsible for this aggravation" he adds.

More information: Detrimental Effects of Diet-Induced Obesity on τ Pathology Is Independent of Insulin Resistance in τ Transgenic Mice, by Antoine Leboucher et al.

Provided by Institut National de la Sante et de la Recherche Medicale

Citation: Detrimental effect of obesity on lesions associated with Alzheimer's disease (2013, January 7) retrieved 20 April 2024 from https://medicalxpress.com/news/2013-01-detrimental-effect-obesity-lesions-alzheimer.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.