

## High anthocyanin intake tied to lower MI risk in younger women

January 15 2013



High intake of a specific sub-class of flavonoids, called anthocyanins, is associated with a reduced risk of myocardial infarction in young and middle-aged women, according to a study published in the Jan. 15 issue of *Circulation*.

(HealthDay)—High intake of a specific sub-class of flavonoids, called anthocyanins, is associated with a reduced risk of myocardial infarction (MI) in young and middle-aged women, according to a study published in the Jan. 15 issue of *Circulation*.

To examine the correlation between anthocyanin and other flavonoid intake and the risk of MI, Aedín Cassidy, Ph.D., from the Norwich Medical School at the University of East Anglia in the United Kingdom, and colleagues followed 93,600 women (25 to 42 years of age) participating in the Nurses' Health Study II who were healthy at baseline (1989). A validated food-frequency questionnaire collected every four years was used to calculate flavonoid intake.



The researchers found that, during 18 years of follow-up, there were 405 cases of MI. After multivariable adjustment, there was a significant inverse association between higher intake of anthocyanins and risk of MI (hazard ratio [HR], 0.68 for highest versus lowest quintiles). The relationship remained significant even after inclusion of intermediate conditions, including history of hypertension (HR, 0.70). Combined intake of blueberries and strawberries correlated with a non-significantly decreased risk of MI (HR, 0.66; 95 percent confidence interval, 0.40 to 1.08) when comparing those consuming more than three servings a week to those with lower intake. Other flavonoid subclasses were not significantly correlated with MI risk.

"Our findings suggest that <u>bioactive compounds</u> present in red and blue <u>fruits and vegetables</u> commonly consumed in the habitual diet may be associated with a reduced risk of MI in young and middle-aged women," the authors write.

**More information:** Abstract

Full Text (subscription or payment may be required)

Copyright © 2013 HealthDay. All rights reserved.

Citation: High anthocyanin intake tied to lower MI risk in younger women (2013, January 15) retrieved 25 April 2024 from

https://medicalxpress.com/news/2013-01-high-anthocyanin-intake-tied-mi.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.