

Migraine triggers may not be as strong as you think

January 23 2013

A new study suggests that triggers for migraine with aura may not be as strong as some people think. The research is published in the January 23, 2013, online issue of *Neurology*, the medical journal of the American Academy of Neurology. Auras that occur with migraine include visual disturbances, with symptoms such as flashing lights or wavy lines.

"People with migraine with aura are told to avoid possible triggers, which may lead them to avoid a wide range of suspected factors," said study author Jes Olesen, MD, with the University of Copenhagen in Denmark and a Fellow of the American Academy of Neurology. "Yet the most commonly reported triggers are stress, bright light, emotional influences and physical effort, which can be difficult to avoid and potentially detrimental, if people avoid all physical activity."

The study involved 27 people with migraine with aura who reported that bright or flickering light, [vigorous exercise](#), or both, previously triggered an attack. The participants were then exposed to the triggers to see if they caused a headache episode.

Participants either went for an intense run or used an [exercise bike](#) for one hour, reaching at least 80 percent of their [maximum heart rate](#). Participants also were exposed to bright, flashing or flickering lights for 30 to 40 minutes. After each session, the participants were monitored for about three hours and asked to report any migraine or migraine with aura symptoms.

The study found that 11 percent of participants reported a migraine with aura after being exposed to light or exercise. Another 11 percent of participants experienced migraines without aura. No participants developed migraine with aura after [light exposure](#) alone.

"Our study suggests that if a person is exposed to a suspected trigger for three months and does not have a [migraine attack](#), they no longer have to worry about avoiding that trigger," said Olesen.

Peter Goadsby, MD, PhD, with the University of California, San Francisco, and a member of the American Academy of Neurology, noted in an accompanying editorial that the study brings up several questions about migraine triggers. "Perhaps rather than triggers, these behaviors are a brain-driven response to the early phases of the migraine itself. Maybe people are driven to exercise as an early symptom and the association with light is simply the sensitivity to light that occurs with the attack itself?"

Provided by American Academy of Neurology

Citation: Migraine triggers may not be as strong as you think (2013, January 23) retrieved 19 April 2024 from <https://medicalxpress.com/news/2013-01-migraine-triggers-strong.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--