

Study finds parasites and poor antenatal care are main causes of epilepsy in Africa

January 30 2013

The largest study of epilepsy in sub-Saharan Africa to date reveals that programmes to control parasitic diseases and access to better antenatal care could substantially reduce the prevalence of the disease in this region.

Epilepsy is one of the most common [neurological conditions](#) worldwide and it is well known that it is significantly more prevalent in poorer countries and rural areas. The study of over half a million people in five countries of sub-Saharan Africa is the first to reveal the true extent of the problem and the impact of different risk factors.

The study, conducted at International Network for the Demographic Evaluation of Populations and Their Health (INDEPTH) demographic surveillance sites in Kenya, South Africa, Uganda, Tanzania and Ghana, screened 586,607 residents and identified 1711 who were diagnosed as having active convulsive epilepsy. These individuals, along with 2033 who did not have epilepsy, were given a questionnaire to complete about their [lifestyle habits](#). The team also took [blood samples](#) to test for exposure to malaria, HIV and four other parasitic diseases that are common in the developing world.

The team found that adults who had been exposed to [parasitic diseases](#) were 1.5-3 times more likely to have epilepsy than those who had not. Epilepsy has previously been linked with various parasite infections but this is the first study to reveal the extent of the problem.

Professor Charles Newton from the Wellcome Trust programme at the Kenyan Medical Research Institute (KEMRI) and the Department of Psychiatry at Oxford University, who led the study, said: "This study demonstrates that many cases of epilepsy could be entirely preventable with elimination of parasites in Africa, some of which for example [onchocerciasis](#) have been controlled in some areas. In some areas the incidence of epilepsy could be reduced by 30-60% with appropriate [control measures](#)."

In children, the greatest [risk factors](#) for developing epilepsy were complications associated with delivery and head injury. Interventions to improve antenatal and perinatal care could substantially reduce the prevalence of epilepsy in this region, say the authors.

The study focused on people with convulsive epilepsies as they are the most reliably detected and reported and there remains a substantial stigma attached to patients with the disease.

"Facilities for diagnosis, treatment and on-going management of epilepsy are virtually non-existent in many of the world's poorest regions and so it's vital that we take these simple steps to try and prevent as many cases of this debilitating disease as possible," Professor Newton added.

More information: A.K. Ngugi et al. Prevalence of active convulsive epilepsy in sub-Saharan Africa and associated risk factors: cross-sectional and case-control studies. *Lancet Neurology*, 31 January 2013.

Provided by Wellcome Trust

Citation: Study finds parasites and poor antenatal care are main causes of epilepsy in Africa

(2013, January 30) retrieved 19 April 2024 from

<https://medicalxpress.com/news/2013-01-parasites-poor-antenatal-main-epilepsy.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.