

Study finds that people with cystic fibrosis who live in deprived areas have worse health

January 29 2013

A study by the University of Liverpool has found that people with Cystic Fibrosis (CF) who live in deprived areas have worse growth and lung function than people living in more advantaged areas.

Researchers at the University of Liverpool examined the medical records of more than 9,000 cystic fibrosis patients under 40 years old, from 1996 to 2009. They found that children and adults with CF who lived in the most disadvantaged areas in the UK had lower weight, height and [Body Mass Index](#) (BMI) than those from more affluent areas.

They were also more likely to have worse lung function, and to pick up severe [lung infections](#). These differences appeared early on in childhood and continued through to adulthood, but did not get any wider over time.

The study, which was funded by the Medical Research Council (MRC) and used data provided by the Cystic Fibrosis Trust, also looked at the treatments received by people with CF in the NHS. It found that the NHS takes this disadvantage into account and treats people from deprived areas more intensively. They were about twice as likely to be treated with [intravenous antibiotics](#) and nutritional therapies, even after taking into account their worse health. However, they were less likely to be treated with inhaled treatments, such as DNase, and inhaled antibiotics.

Dr David Taylor-Robinson, from the Department of Public Health and Policy, said: "This is the first study to examine the relationship between

deprivation and longitudinal outcomes in [cystic fibrosis](#) in the UK, which is of particular interest in the study of [health inequalities](#) because it is a genetic disease and affects all [socioeconomic groups](#) equally.

"It is important that children with CF are not exposed to second-hand smoke, which is more common in deprived areas, and may explain some of the difference in early lung function. This study also shows that the health effects of deprivation appear very early in childhood, even for [genetic diseases](#) such as CF.

"We need to make sure that all children get the best possible start in life in order to reduce health inequalities."

The research is published in *The Lancet Respiratory Medicine*.

Provided by University of Liverpool

Citation: Study finds that people with cystic fibrosis who live in deprived areas have worse health (2013, January 29) retrieved 27 April 2024 from <https://medicalxpress.com/news/2013-01-people-cystic-fibrosis-deprived-areas.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--