

Plumber and spray painter high-risk occupations for asthma

January 15 2013

Despite known risks and recommendations for protective equipment, many people are still affected with asthma after exposure to chemicals at work. This is the finding of an international study of 13,000 people carried out at the Sahlgrenska Academy, University of Gothenburg, Sweden.

Asthma is among the most common adult diseases in the world. Despite the fact that the risks of [chemical exposure](#) have long been known and that there are well-established recommendations for handling chemicals and [protective equipment](#), many cases of asthma are still caused by exposure to toxic substances at work.

A study at Sahlgrenska Academy, University of Gothenburg, has tracked asthma cases among 13,000 randomly selected adults in Sweden, Norway, Denmark, Iceland and Estonia from 1980 to 2000. According to the study, 429 people had asthma during this period. Seven percent of cases among women were linked to workplace exposure—and among men, the number

The study found that total incidence was 1.3 asthma cases per 1,000 men, and 2.4 cases per 1,000 women. "To be able to work proactively, it is essential to show which substances at work increase the risk of asthma and which occupations are high-risk," says Linnea Lillienberg, researcher at Sahlgrenska Academy, University of Gothenburg.

According to the study, high-risk occupations include:

- spray painters, who are exposed to diisocyanates in paint
- plumbers, who handle adhesives and [foam insulation](#)
- cleaners, who handle [detergents](#)
- health care and social services personnel, who are exposed to detergents and latex in latex gloves, especially if the gloves contain powder
- food and [tobacco industry](#) workers, who are exposed to proteins from the vegetable kingdom
- hair stylists, who handle chemicals in bleach and nail beauticians, who use fast-acting glue.

"Some people are more susceptible than others. For example, people with [hay fever](#) have asthma more often if they're exposed to proteins from [plants and animals](#). But if we look at individuals with no increased susceptibility, the risk was greater among those who were exposed to epoxy and diisocyanates, which are found in glue, varnish and foam plastic. Among women without hay fever, the risk was particularly elevated among those who handled detergents," says Linnea Lillienberg.

The study, "Occupational Exposure and New-onset Asthma in a Population-based Study in Northern Europe (RHINE)" was published in *The Annals of Occupational Hygiene*.

Provided by University of Gothenburg

Citation: Plumber and spray painter high-risk occupations for asthma (2013, January 15) retrieved 23 April 2024 from <https://medicalxpress.com/news/2013-01-plumber-painter-high-risk-occupations-asthma.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.