

# Study finds poorer outcomes for obese patients treated for lumbar disc herniation

January 10 2013

---

While obese patients are more likely to have surgical treatment for lumbar disc herniation – a slipped or ruptured disc – than nonobese patients, obesity increases operative time, blood loss and length of hospital stay, according to new research published in the January 2013 *Journal of Bone and Joint Surgery* (JBJS). Overall, obese patients had poorer outcomes with surgical and nonsurgical treatments for lumbar disc herniation than nonobese patients.

The study included 854 nonobese patients with a [Body Mass Index](#) (BMI) of less than 30 kg/m<sup>2</sup>, and 336 [obese patients](#) with a BMI greater than 30 kg/m<sup>2</sup>, enrolled in the Spine Patient Outcomes Research Trial (SPORT) for the treatment of lumbar disc herniation. Researchers compiled patient demographic and clinical characteristics at baseline, and then compared that information with data compiled during regular follow-up visits for four years.

At four years, improvements in function were less for obese patients – in both the surgical and nonsurgical groups – as measured by the [Oswestry Disability Index](#). Reported pain (using the Short Form-36 scale) was statistically similar for obese and non-obese patients.

"The findings suggest that obese patients with symptomatic lumbar disc herniation do not do as well as nonobese patients with nonsurgical or surgical treatment," said Jeffrey Rihn, MD, associate professor at Thomas Jefferson University Hospital and The Rothman Institute.

## Other Key Findings:

- Obese patients did not have an increased rate of infection, intraoperative complications, recurrent disc herniation or reoperation.
- Obese patients had significantly less improvement in the Sciatica Bothersomeness Index and [Low Back Pain](#) Bothersomeness Index, but reported no significant difference in self-rated overall improvement/satisfaction.
- The benefit of surgery over nonoperative treatment was not affected by body mass index.
- Recurrence of disc herniation and need for additional surgical procedures did not differ significantly between obese and nonobese patients. This finding contradicts previous studies on this topic.

"The results of this study may be helpful in educating patients about their treatment options and expected outcomes," said Dr. Rihn. "These findings may suggest that weight loss should be encouraged in patients with this condition. However, this study does not specifically address whether weight loss in obese patients would affect their clinical outcome with nonsurgical or surgical treatment."

Provided by American Academy of Orthopaedic Surgeons

Citation: Study finds poorer outcomes for obese patients treated for lumbar disc herniation (2013, January 10) retrieved 5 May 2024 from <https://medicalxpress.com/news/2013-01-poorer-outcomes-obese-patients-lumbar.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
---