

# Surgery consultation common after MRI of the spine

January 2 2013

---



Almost half of patients whose primary care physicians recommend a lumbosacral or cervical spine magnetic resonance imaging scan go on to receive a surgical consultation, but few end up undergoing spinal surgery, according to research published in the Jan. 1 issue of *Spine*.

(HealthDay)—Almost half of patients whose primary care physicians recommend a lumbosacral or cervical spine magnetic resonance imaging (MRI) scan go on to receive a surgical consultation, but few end up undergoing spinal surgery, according to research published in the Jan. 1 issue of *Spine*.

John J. You, M.D., of McMaster University in Hamilton, Canada, and colleagues evaluated subsequent patterns of health care utilization in a [retrospective cohort study](#) involving 647 patients who had a lumbosacral MRI and 373 patients who had a cervical spine MRI scan ordered by a primary care physician.

After lumbosacral spine MRI, the researchers found that 44.5 percent of patients were seen by an orthopedic surgeon or neurosurgeon, and 6.5 percent ultimately underwent spine surgery during the subsequent three years. After cervical spine MRI, 44.0 percent were seen by an orthopedic surgeon or neurosurgeon, and none had surgery. Subsequent surgery was more likely for patients with severe disc herniation (likelihood ratio, 5.62) or severe spinal stenosis (likelihood ratio, 2.34) on lumbosacral spine MRI. Not all patients with these abnormalities underwent surgery and the absence of these findings was not associated with a significantly lower likelihood of subsequent surgery.

"We determined that [MRI scans](#) of the spine performed in symptomatic patients in primary care were almost always abnormal, subsequent surgical referral was frequent, and subsequent surgery was infrequent," the authors write. "Results of MRI scans were not strongly predictive of subsequent surgery and underscore the importance of clinical features of pain, functional limitation, and neurological involvement, and consideration of individual patients' values and preferences when making decisions about spinal surgery."

One or more authors disclosed receipt of a benefit from a commercial party.

**More information:** [Abstract](#)  
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2012 [HealthDay](#). All rights reserved.

Citation: Surgery consultation common after MRI of the spine (2013, January 2) retrieved 2 May 2024 from <https://medicalxpress.com/news/2013-01-surgery-common-mri-spine.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.