

Vaccination responsible for dramatic fall in salmonella infections

January 16 2013



Mass poultry vaccination programmes introduced to combat Salmonella infections have led to a dramatic fall in the number of cases.

Mass poultry vaccination programmes introduced to combat Salmonella infections have led to a dramatic fall in the number of cases since the late 1990s, according to a researcher at the University of Liverpool.

[Salmonella](#) are important food-borne pathogens worldwide, causing diarrhoea, vomiting, nausea, fever and abdominal pain. There are currently around 6 million cases of illness from Salmonella across the

EU each year, the majority of which are linked to food items such as eggs, chicken, beef, pork, salad vegetables and dairy products.

Between 1981 and 1991, the number of [salmonella infections](#) rose by 170% in the UK, driven primarily by an epidemic of Salmonella Enteritidis which peaked in 1993. A raft of [control measures](#) were introduced into the [poultry industry](#) including movement restrictions, compulsory slaughter and disinfection procedures, as well as a voluntary industry-led vaccination scheme that began in breeding [flocks](#) in 1994 and in laying flocks in 1998.

Code of practice

Legislation requiring compulsory slaughter of poultry infected with Salmonella has now been revoked but the [mass vaccination](#) of poultry has continued by those breeders subscribing to the Lion Quality Code of Practice and using the Lion Mark on eggs. The code of practice requires [mandatory vaccination](#) of all young hens destined to lay Lion eggs against Salmonella, as well as traceability of hens, eggs and feed, a best-before date stamped on shells and hygiene controls at packing stations. Lion eggs now account for around 85% of the total market.

Sarah O'Brien, Professor of Epidemiology and Zoonoses, from the University's Institute of Infection and [Global Health](#), attributes a dramatic fall in the number of Salmonella cases in humans to this mass vaccination programme in poultry.

Professor O'Brien said: "We have seen a marked decline in the number of incidences of Salmonella infection, shown by two significant studies conducted 10 years apart. These studies found that the number of cases fell from 1.6 cases per 1,000 person years for a study conducted from 1993 to 1996 to 0.2 cases per 1,000 person years for the same study conducted from 2008 to 2009.

"In addition, the number of laboratory-confirmed cases of illness dropped from more than 18,000 in 1993 to just 459 in 2010.

"The nature of public health interventions often means that evaluating their impact is complex as they are often implemented simultaneously. The decrease in laboratory confirmed human cases coincides quite closely with the introduction of vaccination programmes in breeder and laying flocks. It is probable that no single measure contributed to the decline in Salmonella cases but the relationship between vaccination programmes and the reduction in human disease is compelling and suggests these programmes have made a major contribution to improving public health."

The research is published in *Clinical Infectious Diseases*.

Provided by University of Liverpool

Citation: Vaccination responsible for dramatic fall in salmonella infections (2013, January 16) retrieved 19 April 2024 from

<https://medicalxpress.com/news/2013-01-vaccination-responsible-fall-salmonella-infections.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--