

Vaginal delivery safest option for women with pelvic girdle pain, study finds

January 17 2013



(Medical Xpress)—Caesarean section increases the risk of persistent pelvic girdle pain after delivery compared with vaginal delivery, according to a new study from the Norwegian Institute of Public Health.

Caesarean section rates are increasing worldwide, and this trend has partly been explained by women's requests for planned caesarean section without a medical reason. Pregnancy-related pelvic girdle pain has been associated with increased preference for caesarean section and with increased planned caesarean section rates.

"Some women with severe pelvic girdle pain might fear that a vaginal delivery will be too difficult or painful, or will worsen the condition after delivery – even though scientific evidence is lacking", says Elisabeth K. Bjelland, who is a [postdoctoral researcher](#) at the Institute's

Division of Mental Health and lead author of the study.

Knowledge about how caesarean section affects prognosis of pelvic girdle pain is therefore needed. The purpose of this study was to investigate the relationship between mode of delivery and the [persistence](#) of pelvic girdle pain six months after delivery.

In this follow-up study, 79.9% of women had unassisted vaginal deliveries, 6.7% had instrumental vaginal deliveries (by vacuum or forceps), 7.2% had emergency caesarean sections and 6.2% had planned caesarean sections.

Doubles the risk of pain after delivery

After taking a number of other factors into account, the results showed that:

- Planned caesarean section was associated with a two- to threefold increased risk of severe pelvic girdle syndrome ([severe pain](#) over the frontal part of the pelvis and on both sides of the rear part of the pelvis) six months after delivery.
- In women who used [crutches](#) in pregnancy, both emergency and planned caesarean section was associated with severe pelvic girdle syndrome six months after delivery.
- Instrumental vaginal delivery was also associated with an increased risk of pelvic girdle syndrome six months after delivery.

"Previous studies have reported that patients with prolonged or severe pain are at increased risk of chronic pain after surgery. One explanation may be that pain influences the central nervous system in such a way that the pain inhibition systems become less effective. Therefore, women

with severe pelvic girdle pain during pregnancy may be more vulnerable to the pain associated with caesarean section, compared with women without pelvic girdle pain", explains Bjelland.

Clinical advice

When deciding the mode of delivery, clinicians should be aware of the increased risk of non-recovery after caesarean section among women with severe pelvic girdle pain.

"Our findings, which support previous assumptions, do not indicate that caesarean section represents a benefit for the process of recovery from pelvic girdle pain. If there are no medical reasons for a [caesarean section](#), the findings suggest that [vaginal delivery](#) is the safest option for women presenting with severe [pelvic girdle pain](#)", advises Bjelland.

More information: Bjelland E.K., Stuge B., Vangen S., Stray-Pedersen, B. and Eberhard-Gran, M. Mode of delivery and persistence of pelvic girdle syndrome 6 months postpartum. *Am J Obstet Gynecol* (2013); 208

Provided by Norwegian Institute of Public Health

Citation: Vaginal delivery safest option for women with pelvic girdle pain, study finds (2013, January 17) retrieved 25 April 2024 from <https://medicalxpress.com/news/2013-01-vaginal-delivery-safest-option-women.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
