

Abnormal growth regulation may occur in children with heart defects

February 19 2013

The poor growth seen in children born with complex heart defects may result from factors beyond deficient nutrition. A new study by pediatric researchers suggests that abnormalities in overall growth regulation play a role.

"When compared with their healthy peers, children with [congenital heart disease](#) have impaired growth, as measured in weight, length, and [head circumference](#)," said senior author Meryl S. Cohen, M.D., a pediatric cardiologist in the Cardiac Center at The Children's Hospital of Philadelphia. "We investigated patterns of poor growth in these children, as a starting point in guiding us toward more effective treatments."

The study appeared as an online article in the January 2013 issue of *Pediatrics*.

The researchers performed a retrospective analysis of medical records of 856 children with congenital heart disease (CHD), compared to 7,654 matched control subjects. All the children were measured up to age 3, and all were drawn from the healthcare network of The Children's Hospital of Philadelphia.

Within weeks of birth, the children with CHD had significant deficits in weight, length and head circumference, compared to matched controls without CHD. The largest differences in weight occurred at 4 months of age. Among the 856 children with CHD, the 248 who required surgical repair were much more likely to be below the 3rd percentile in weight,

length and head circumference during early [infancy](#), and their growth by age 3 did not catch up with that of their healthy peers.

In the 608 children with CHD who did not require surgery, growth differences were not as pronounced, but even their growth patterns lagged behind those of healthy controls.

Researchers already knew that children with CHD have an increased risk for poor growth, but this analysis provides a fuller picture of the problem. Cohen observed that in the general population, when [caloric intake](#) is insufficient, an infant's weight is usually affected first, followed by length and head circumference. "The fact that all three parameters changed simultaneously rather than sequentially supports the idea that impaired growth in children with heart disease is affected at least in part by factors unrelated to nutrition."

She added that further studies should investigate the possible roles of growth hormones and other physiologic factors that affect growth regulation in children with CHD.

More information: *Pediatrics*, Jan. 2013, pp. e236-e242

Provided by Children's Hospital of Philadelphia

Citation: Abnormal growth regulation may occur in children with heart defects (2013, February 19) retrieved 12 May 2024 from

<https://medicalxpress.com/news/2013-02-abnormal-growth-children-heart-defects.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--