

Amyloid imaging shows promise for detecting cardiac amyloidosis

February 4 2013

While amyloid imaging may now be most associated with detecting plaques in the brain, it has the potential to change the way cardiac amyloidosis is diagnosed. According to first-of-its-kind research published in the February issue of *The Journal of Nuclear Medicine*, positron emission tomography (PET) with ^{11}C -PIB can positively visualize amyloid deposits in the heart. Currently there is no noninvasive test available for specific diagnosis.

Cardiac amyloidosis is a deadly disorder caused by abnormal amyloid deposits in the heart tissue. Early diagnosis before structural change to the [heart tissue](#) has occurred is important for disease prognosis and for treatment monitoring. Currently echocardiography is the mainstay of imaging in cardiac amyloidosis; a cardiac biopsy is used to confirm diagnosis.

"Imaging with ^{11}C -PIB provides a non-invasive and specific means of showing distribution of amyloid in an organ. This gives a unique opportunity to follow and monitor therapy, as amyloid deposits in the heart should decrease with successful therapy," said Gunnar Antoni, PhD, lead author of the study "In Vivo Visualization of Amyloid Deposits in the Heart with ^{11}C -PIB and PET."

The study included 10 patients diagnosed with cardiac amyloidosis and five healthy individuals. PET/computed tomography (CT) with ^{11}C -PIB was used to visualize [amyloid deposits](#) in the heart and with ^{11}C -acetate to measure myocardial blood flow.

Uptake of ^{11}C -PIB was measured 15-25 minutes after injection. Obvious uptake of ^{11}C -PIB was noted in the [left ventricle](#) wall of all patients with cardiac amyloidosis, while no uptake was seen in the healthy volunteers. In half of the patients, ^{11}C -PIB was also detected in the [right ventricle](#) wall, and nine of the patients had signs of reversible uptake, with a maximum concentration at 10-15 minutes after injection. Myocardial blood flow was significantly lower in patients with cardiac amyloidosis; however, no significant correlation between myocardial blood flow and ^{11}C -PIB uptake was found.

"This study emphasizes the strength of molecular imaging for detecting an underlying and significant molecular aberration in a disease that presents with unspecific symptoms and signs," noted Antoni. "The potential for molecular imaging to provide valuable information for other diseases is of great value to the field of medicine."

More information: "In Vivo Visualization of Amyloid Deposits in the Heart with ^{11}C -PIB and PET", *The Journal of Nuclear Medicine*, 2013.

Provided by Society of Nuclear Medicine

Citation: Amyloid imaging shows promise for detecting cardiac amyloidosis (2013, February 4) retrieved 8 May 2024 from <https://medicalxpress.com/news/2013-02-amyloid-imaging-cardiac-amyloidosis.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--