

## One in 20 cases of pre-eclampsia may be linked to air pollutant

February 6 2013

---

One in every 20 cases of the serious condition of pregnancy, pre-eclampsia, may be linked to increased levels of the air pollutant ozone during the first three months, suggests a large study published in the online journal *BMJ Open*.

Mothers with asthma may be more vulnerable, the findings indicate.

Pre-eclampsia is characterised by raised blood pressure and the presence of protein in the urine during pregnancy. It can cause serious complications, if left untreated.

The authors base their findings on almost 121,000 [singleton births](#) in Greater Stockholm, Sweden, between 1998 and 2006; national data on the prevalence of asthma among the children's mothers; and levels of the [air pollutants](#) ozone and vehicle exhaust (nitrogen oxide) in the Stockholm area.

There's a growing body of evidence pointing to a link between air pollution and premature birth, say the authors, while pregnant women with asthma are more likely to have [pregnancy complications](#), including underweight babies and pre-eclampsia.

In all, 4.4% of the pregnancies resulted in a premature birth and the prevalence of pre-eclampsia was 2.7%.

There was no association between exposure to levels of [vehicle exhaust](#)

and complications of pregnancy, nor were any associations found for any air pollutants and babies that were underweight at birth.

But there did seem to be a link between exposure to ozone levels during the first three months of pregnancy and the risk of premature birth (delivery before 37 weeks) and pre-eclampsia, after adjusting for factors likely to influence the results and [seasonal variations](#) in air pollutants, although not spatial variations in exposure.

Each rose by 4% for every 10 ug/m<sup>3</sup> rise in ambient ozone during this period, the analysis indicated.

Mothers with asthma were 25% more likely to have a child born prematurely and 10% more likely to have pre-eclampsia than mums without this condition.

Asthma is an inflammatory condition and ozone may therefore have worsened respiratory symptoms and systemic inflammation, so accounting for the larger increase in the risk of premature birth among the mums with asthma, suggest the authors.

But after taking account of the mother's age, previous births, educational attainment, ethnicity, asthma, season and year of conception, the authors calculated that one in every 20 (5%) cases of pre-eclampsia were linked to [ozone levels](#) during early pregnancy.

**More information:** [www.bmjopen.bmj.com/lookup/doi ... /bmjopen-2012-001955](http://www.bmjopen.bmj.com/lookup/doi.../bmjopen-2012-001955)

Provided by British Medical Journal

Citation: One in 20 cases of pre-eclampsia may be linked to air pollutant (2013, February 6)  
retrieved 25 April 2024 from

<https://medicalxpress.com/news/2013-02-cases-pre-eclampsia-linked-air-pollutant.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.