

# The criteria for weight-loss surgery need to be changed

February 19 2013

---

(Medical Xpress)—Weight-loss surgery is currently only offered to patients who exceed a certain BMI. However, surgical intervention could improve the health of many more people. This is shown by the Swedish Obese Subjects study carried out at the Sahlgrenska Academy, University of Gothenburg, Sweden, involving 104 patients who were operated on despite their BMI being "too low". As a result, the risk of developing diabetes was reduced by 67 percent.

In order to meet the current selection criteria for weight-loss surgery, Swedish patients must have a [body mass index](#) (BMI) above 40 for healthy patients, or above 35 in combination with other risk factors such as [type 2 diabetes](#).

However, these selection criteria have not been devised in a scientific manner, and need to be improved. This is the opinion of researchers at the Sahlgrenska Academy, University of Gothenburg, who have shown in a new study that weight-loss surgery also offers significant health benefits for patients who do not meet today's criteria.

In the study, which has been published in [Diabetes Care](#), the researchers from Gothenburg have monitored 104 patients who had weight-loss surgery before the current criteria were established, and who would not have been offered the operation today. The study shows that, 15 years after the procedure, these patients had a 67 percent lower risk of developing [diabetes](#) compared with a control group who were not operated on.

"The effect was the same as for patients who meet the current criteria," explains researcher Kajsa Sjöholm from the Sahlgrenska Academy, University of Gothenburg. "In other words, there is no proof that weight-loss surgery has a greater preventive effect against diabetes for patients who exceed today's BMI limit than for those who don't."

The study also shows that weight-loss surgery reduces the risk factors for cardiovascular disease among patients who are not currently eligible for this surgery.

With reference to previous results published in the SOS study at the University of Gothenburg, which showed that weight-loss surgery generally results in a long-lasting reduction in weight and reduces the risk of diabetes, heart disease, stroke, cancer and premature death, the researchers from Gothenburg believe that today's selection criteria should be re-evaluated.

According to the researchers, strict BMI limits can lead to patients with a high risk of future disease who would benefit from treatment being denied [surgery](#). Previous research findings show that the selection can be improved by analysing patients' insulin and glucose values.

Provided by University of Gothenburg

Citation: The criteria for weight-loss surgery need to be changed (2013, February 19) retrieved 27 April 2024 from <https://medicalxpress.com/news/2013-02-criteria-weight-loss-surgery.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--