

Rewriting a receptor's role: Synaptic molecule works differently than thought

February 19 2013

(Medical Xpress)—In a pair of new papers, researchers at the University of California, San Diego School of Medicine and the Royal Netherlands Academy of Arts and Sciences upend a long-held view about the basic functioning of a key receptor molecule involved in signaling between neurons, and describe how a compound linked to Alzheimer's disease impacts that receptor and weakens synaptic connections between brain cells.

The findings are published in the Feb. 18 early edition of the *Proceedings of the National Academy of Sciences*.

Long the object of study, the NMDA receptor is located at [neuronal synapses](#) – the multitudinous junctions where brain cells trade electrical and chemical messages. In particular, NMDA receptors are ion channels activated by glutamate, a major "excitatory" neurotransmitter associated with cognition, learning and memory.

"NMDA receptors are well known to allow the passage of calcium ions into cells and thereby trigger biochemical signaling," said principal investigator Roberto Malinow, MD, PhD professor of neurosciences at UC San Diego School of Medicine.

The new research, however, indicates that NMDA receptors can also operate independent of [calcium ions](#). "It turns upside down a view held for decades regarding how NMDA receptors function," said Malinow, who holds the Shiley-Marcos Endowed Chair in Alzheimer's Disease

Research in Honor of Dr. Leon Thal (a renowned UC San Diego Alzheimer's disease researcher who died in a single-engine [airplane crash](#) in 2007).

Specifically, Malinow and colleagues found that glutamate binding to the NMDA receptor caused conformational changes in the receptor that ultimately resulted in a weakened synapse and impaired brain function.

They also found that beta amyloid – a peptide that comprises the neuron-killing plaques associated with Alzheimer's disease – causes the [NMDA receptor](#) to undergo conformational changes that also lead to the weakening of synapses.

"These new findings overturn commonly held views regarding synapses and potentially identify new targets in the treatment of Alzheimer's disease," said Malinow.

Provided by University of California - San Diego

Citation: Rewriting a receptor's role: Synaptic molecule works differently than thought (2013, February 19) retrieved 3 May 2024 from <https://medicalxpress.com/news/2013-02-rewriting-receptor-role-synaptic-molecule.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--