

# Strains of antibiotic-resistant 'Staph' bacteria show seasonal preference: Children at higher risk in summer

February 28 2013

---

Strains of potentially deadly, antibiotic-resistant *Staphylococcus aureus* bacteria show seasonal infection preferences, putting children at greater risk in summer and seniors at greater risk in winter, according to results of a new nationwide study led by a Johns Hopkins researcher.

It's unclear why these seasonal and age preferences for infection with methicillin-resistant *Staph aureus* (MRSA) occur, says Eili Klein, Ph.D., lead author on the study and a researcher at the Johns Hopkins Center for Advanced Modeling in the Social, Behavioral and [Health Sciences](#).

But he says that increased [use of antibiotics](#) in the winter may be one of the reasons. The winter strain that infects seniors at a greater rate is generally acquired in the hospital and resistant to more antibiotics. On the other hand, the summer strain of MRSA, which is seen with growing frequency in children, is largely a community-transmitted strain that is resistant to fewer antibiotics.

"Overprescribing antibiotics is not harmless," Klein notes.

"Inappropriate use of these drugs to treat [influenza](#) and other respiratory infections is driving resistance throughout the community, increasing the probability that children will contract untreatable infections."

In fact, the study found that while MRSA [strains](#) exhibit a [seasonal pattern](#), overall MRSA infections have not decreased over the last five

years, despite efforts to control their spread.

A report on the study, which used sophisticated statistical models to analyze national data for 2005-2009, appears today in the online issue of the [American Journal of Epidemiology](#).

As the researchers report, hospitalizations from infections tied to MRSA doubled in the United States between 1999 and 2005. The ballooning infection numbers were propelled by MRSA acquired in community settings, not hospital or other health care settings, as had been the case prior to 1999.

Specifically, the study found that a strain of MRSA typically seen in community settings is more likely to cause infection during the summer months, peaking around July/August. The authors' data analysis showed children were most at risk of becoming infected with this strain, typically from a skin or soft tissue wound or ailment.

In fact, in examining data for one year—2008—the research team found that 74 percent of those under the age of 20 who developed an infection with MRSA had a community-associated MRSA infection.

Meanwhile, the health care-associated MRSA strain, which is typically seen in hospitals, nursing homes and other health care settings, was found to be most prevalent in the winter months, peaking in February/March. Patients aged 65 or older are more likely to acquire a MRSA infection from this strain.

"Our analysis ... shows significant seasonality of MRSA infections and the rate at which they affect different age groups," write the authors of the report titled "The changing epidemiology of methicillin-resistant *Staphylococcus aureus* in the United States: A national observational study."

Klein said additional research on seasonal patterns of [MRSA infections](#) and drug resistance may help with developing new treatment guidelines, prescription practices and infection control programs.

**More information:** [aje.oxfordjournals.org/content ...  
/aje.kws273.abstract](http://aje.oxfordjournals.org/content.../aje.kws273.abstract)

Provided by Johns Hopkins University School of Medicine

Citation: Strains of antibiotic-resistant 'Staph' bacteria show seasonal preference: Children at higher risk in summer (2013, February 28) retrieved 9 April 2024 from <https://medicalxpress.com/news/2013-02-strains-antibiotic-resistant-staph-bacteria-seasonal.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--