

Strokes associated with surgery can be devastating

February 11 2013

Strokes that occur during or shortly after surgery can be devastating, resulting in longer hospital stays and increased risks of death or long-term disability.

But prompt identification and treatment of such strokes can improve neurologic outcomes, according to an article in the journal *Expert Review of [Neurotherapeutics](#)* by Loyola University Medical Center stroke specialists Sarkis Morales-Vidal, MD and Michael Schneck, MD.

The article answers commonly asked questions about the management of perioperative stroke. (A perioperative stroke is a stroke that occurs anytime between the time a patient is hospitalized for surgery until the patient is discharged from the hospital.)

Risk factors for peroperative stroke include advanced age, female gender, obesity, high blood pressure, smoking, [peripheral vascular disease](#), [chronic obstructive pulmonary disease](#), diabetes and high cholesterol. For most surgeries, the risk of perioperative stroke is less than 1 percent. But the risk can be as high as 5 percent for surgeries for head and neck tumors and between 2 and 10 percent for various heart surgeries.

The most common cause of perioperative stroke is blood clots. [Blood thinners](#) can reduce the risk of strokes, but can increase the risk of bleeding. Morales and Schneck write that in managing surgery patients, physicians must balance the risk of stroke versus the risk of significant

bleeding complications. Studies have found that for many surgeries, including cardiovascular procedures, the benefits of giving patients aspirin (a blood thinner) outweigh the risks of bleeding.

The authors examine the evidence for several therapies to treat perioperative strokes caused by blood clots:

- Intravenous clot-busting drug (rtPA). Because of the risk of bleeding, rtPA is not indicated for patients who have undergone major surgery within the previous 14 days. But rtPA probably is safe following minor surgeries such as [muscle biopsies](#) and dental procedures.

Delivering rtPA by catheter. In this procedure, a high concentration of the clot-busting drug is delivered by a catheter directly to the clot. But using this technique in any patient population, surgical or otherwise, "is not currently substantiated by randomized controlled trials," the authors write.

Provided by Loyola University Health System

Citation: Strokes associated with surgery can be devastating (2013, February 11) retrieved 4 May 2024 from <https://medicalxpress.com/news/2013-02-surgery-devastating.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
